

ARG63288 anti-NALP3 / Cryopyrin antibody

Package: 50 µg
Store at: -20°C

Summary

Product Description	Goat Polyclonal antibody recognizes NALP3 / Cryopyrin
Tested Reactivity	Hu
Predict Reactivity	Cow, Rat
Tested Application	FACS, ICC/IF
Specificity	Variants (NP_001073289.1; NP_004886.3) encode the same isoform.
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	NALP3 / Cryopyrin
Species	Human
Immunogen	C-EKPELTVVFEPSW
Conjugation	Un-conjugated
Alternate Names	NLRP3; CIAS1; FCU; MWS; FCAS; NALP3; C1orf7; PYPAF1; AII/AVP; AGTAVPRL; CRYOPYRIN; cold autoinflammatory syndrome 1; cryopyrin; Muckle-Wells syndrome; chromosome 1 open reading frame 7; PYRIN-containing APAF1-like protein 1; neuronal apoptosis inhibitor protein 3; angiotensin/vasopressin receptor AII/AVP-like; NLR family; pyrin domain containing 3; AII; AVP; CLR1.1; FLJ95925; AII/AVP receptor-like; NACHT domain-; leucine-rich repeat-; and PYD-containing protein 3; NACHT; LRR and PYD containing protein 3; PYRIN-containing APAF1-like protein 1; nucleotide-binding oligomerization domain; leucine rich repeat and pyrin domain containing 3

Application Instructions

Application table	Application	Dilution
	FACS	10 µg/ml
	ICC/IF	10 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


Properties

Form	Liquid
Purification	Purified from goat serum by antigen affinity chromatography.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA

Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.


Bioinformation

Gene Symbol	NLRP3
Gene Full Name	NLR family pyrin domain containing 3
Background	NALP3 gene encodes a pyrin-like protein containing a pyrin domain, a nucleotide-binding site (NBS) domain, and a leucine-rich repeat (LRR) motif. This protein interacts with the apoptosis-associated speck-like protein PYCARD/ASC, which contains a caspase recruitment domain, and is a member of the NALP3 inflammasome complex. This complex functions as an upstream activator of NF-kappaB signaling, and it plays a role in the regulation of inflammation, the immune response, and apoptosis. Mutations in this gene are associated with familial cold autoinflammatory syndrome (FCAS), Muckle-Wells syndrome (MWS), chronic infantile neurological cutaneous and articular (CINCA) syndrome, and neonatal-onset multisystem inflammatory disease (NOMID). Multiple alternatively spliced transcript variants encoding distinct isoforms have been identified for this gene. Alternative 5' UTR structures are suggested by available data; however, insufficient evidence is available to determine if all of the represented 5' UTR splice patterns are biologically valid. [provided by RefSeq, Oct 2008]
Function	NALP3: As the sensor component of the NLRP3 inflammasome, plays a crucial role in innate immunity and inflammation. In response to pathogens and other damage-associated signals, initiates the formation of the inflammasome polymeric complex, made of NLRP3, PYCARD and CASP1 (and possibly CASP4 and CASP5). Recruitment of proCASP1 to the inflammasome promotes its activation and CASP1-catalyzed IL1B and IL18 maturation and secretion in the extracellular milieu (PubMed:28847925). Activation of NLRP3 inflammasome is also required for HMGB1 secretion (PubMed:22801494). The active cytokines and HMGB1 stimulate inflammatory responses. Inflammasomes can also induce pyroptosis, an inflammatory form of programmed cell death. Under resting conditions, NLRP3 is autoinhibited. NLRP3 activation stimuli include extracellular ATP, reactive oxygen species, K(+) efflux, crystals of monosodium urate or cholesterol, amyloid-beta fibers, environmental or industrial particles and nanoparticles, cytosolic dsRNA, etc. However, it is unclear what constitutes the direct NLRP3 activator. Activation in presence of cytosolic dsRNA is mediated by DHX33 (PubMed:23871209). Independently of inflammasome activation, regulates the differentiation of T helper 2 (Th2) cells and has a role in Th2 cell-dependent asthma and tumor growth. During Th2 differentiation, required for optimal IRF4 binding to IL4 promoter and for IRF4-dependent IL4 transcription. Binds to the consensus DNA sequence 5'-GRRGGNRGAG-3'. May also participate in the transcription of IL5, IL13, GATA3, CCR3, CCR4 and MAF. [UniProt]
Highlight	Related products: NALP3 antibodies ; NALP3 ELISA Kits ; NALP3 Duos / Panels ; Anti-Goat IgG secondary antibodies ; Related news: Exploring Antiviral Immune Response RIP1 activation and pathogenesis of NASH
Research Area	NLRP3 Inflammasome Study antibody
Calculated Mw	118 kDa



ARG63288 anti-NALP3 / Cryopyrin antibody ICC/IF image

Immunofluorescence: Paraformaldehyde-fixed U937 cells permeabilized with 0.15% Triton. Cells were stained with ARG63288 anti-NALP3 / Cryopyrin antibody (green) at 10 µg/ml dilution for 1 hour. DAPI (blue) for nuclear staining. Negative control: Unimmunized Goat IgG (green) at 10 µg/ml dilution.



ARG63288 anti-NALP3 / Cryopyrin antibody FACS image

Flow Cytometry: Paraformaldehyde-fixed U937 cells, permeabilized with 0.5% Triton. Cells were stained with ARG63288 anti-NALP3 / Cryopyrin antibody (blue line) at 10 µg/ml dilution for 1 hour, followed by Alexa Fluor® 488 labelled secondary antibody. IgG control: Unimmunized Goat IgG (black line), followed by Alexa Fluor® 488 labelled secondary antibody.