

## ARG63510 anti-E7EA / UBXD8 antibody

Package: 100 µg  
Store at: -20°C

### Summary

Product Description	Goat Polyclonal antibody recognizes E7EA / UBXD8
Tested Reactivity	Hu, Ms
Predict Reactivity	Cow, Rat, Dog
Tested Application	WB
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	E7EA / UBXD8
Species	Human
Immunogen	C-HTEVLFVQDLTDE
Conjugation	Un-conjugated
Alternate Names	UBXD8; Protein E7EA; FAS-associated factor 2; UB3N3B; UB3 domain-containing protein 3B; UB3 domain-containing protein 8; E7EA

### Application Instructions

Application table	Application	Dilution
	WB	0.3 - 1 µg/ml
Application Note	WB: Recommend incubate at RT for 1h. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties

Form	Liquid
Purification	Purified from goat serum by antigen affinity chromatography.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Database links	<a href="#">GeneID: 23197 Human</a> <a href="#">GeneID: 76577 Mouse</a> <a href="#">Swiss-port # Q3TDN2 Mouse</a> <a href="#">Swiss-port # Q96CS3 Human</a>
Background	The protein encoded by this gene is highly expressed in peripheral blood of patients with atopic dermatitis (AD), compared to normal individuals. It may play a role in regulating the resistance to apoptosis that is observed in T cells and eosinophils of AD patients. [provided by RefSeq, Jul 2008]
Research Area	Immune System antibody
Calculated Mw	53 kDa

Images



ARG63510 anti-E TEA / UB XD8 antibody WB image

Western Blot: Jurkat lysate (RIPA buffer, 35 µg total protein per lane) stained with ARG63510 anti-E TEA / UB XD8 antibody at 0.3 µg/ml dilution.



ARG63510 anti-E TEA / UB XD8 antibody WB image

Western blot: 35 µg of Human thymus (A) and Mouse thymus (B) lysates (in RIPA buffer) stained with ARG63510 anti-E TEA / UB XD8 antibody at 1 µg/ml dilution and incubated at RT for 1 hour.