

ARG64061 anti-SNAIL antibody

Package: 100 µg, 50 µg
Store at: -20°C

Summary

Product Description	Goat Polyclonal antibody recognizes SNAIL
Tested Reactivity	Hu, Rat
Predict Reactivity	Ms
Tested Application	IHC-P, WB
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	SNAIL
Species	Human
Immunogen	RKPSDPNRKPNY-C
Conjugation	Un-conjugated
Alternate Names	SNAH; SNAIL; SNA; dJ710H13.1; Protein sna; Protein snail homolog 1; Zinc finger protein SNAI1; SLUGH2; SNAI1

Application Instructions

Application table	Application	Dilution
	IHC-P	5 µg/ml
	WB	0.1 - 1 µg/ml
Application Note	WB: Recommend incubate at RT for 1h. IHC-P: Antigen Retrieval: Steam tissue section in Citrate buffer (pH 6.0). * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Purified from goat serum by antigen affinity chromatography.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links

[GeneID: 6615 Human](#)

[Swiss-port # O95863 Human](#)

Background

The Drosophila embryonic protein snail is a zinc finger transcriptional repressor which downregulates the expression of ectodermal genes within the mesoderm. The nuclear protein encoded by this gene is structurally similar to the Drosophila snail protein, and is also thought to be critical for mesoderm formation in the developing embryo. At least two variants of a similar processed pseudogene have been found on chromosome 2. [provided by RefSeq, Jul 2008]

Research Area

Cancer antibody; Cell Biology and Cellular Response antibody; Controls and Markers antibody; Developmental Biology antibody; Gene Regulation antibody; Neuroscience antibody


Calculated Mw

29 kDa

PTM


Phosphorylated by GSK3B. Once phosphorylated, it becomes a target for BTRC ubiquitination. Phosphorylation by CSNK1E, probably at Ser-104, provides the priming site for the subsequent phosphorylation by GSK3B, probably at Ser-100 and Ser-96. Phosphorylation by PAK1 may modulate its transcriptional activity by promoting increased accumulation in the nucleus. Phosphorylation at Ser-11 and Ser-92 positively regulates its functions in induction of EMT and cell survival, respectively. Phosphorylation by LATS2, upon mitotic stress, oncogenic stress or Hippo pathway activation, occurs in the nucleus and promotes nuclear retention and stabilization of total cellular protein level. Ubiquitinated on Lys-98, Lys-137 and Lys-146 by FBXL14 and BTRC leading to degradation. BTRC-triggered ubiquitination requires previous GSK3B-mediated SNAI1 phosphorylation. Ubiquitination induced upon interaction with NOTCH1 or TP53/p53 is mediated by MDM2. O-GlcNAcylation at Ser-112 is enhanced in hyperglycaemic conditions, it opposes phosphorylation by GSK3B, and stabilizes the protein. ADP-ribosylation by PARP1 increases protein half-life and may be involved in TGF β -induced SNAI1 up-regulation.

Images




ARG64061 anti-SNAI1 antibody WB image

Western blot: 20 μ g of HeLa and A549 cell lysates stained with ARG64061 anti-SNAI1 antibody at 1:1000 dilution.




ARG64061 anti-SNAIL antibody IHC-P image

Immunohistochemistry: Paraffin-embedded breast carcinoma tissue stained with ARG64061 anti-SNAIL antibody at 1:100 dilution. Antigen Retrieval: Boil tissue section in EDTA buffer (pH 9.0).




ARG64061 anti-SNAIL antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human kidney tissue. Antigen Retrieval: Steam tissue section in Citrate buffer (pH 6.0). The tissue section was stained with ARG64061 anti-SNAIL antibody at 5 µg/ml dilution followed by AP-staining.



ARG64061 anti-SNAIL antibody WB image

Western blot: 35 µg of Rat kidney lysate (in RIPA buffer) stained with ARG64061 anti-SNAIL antibody at 0.1 µg/ml dilution and incubated at RT for 1 hour.



ARG64061 anti-SNAIL antibody IHC-P image

Immunohistochemistry: Paraffin-embedded lung tissue from HGF overexpressing mouse stained with ARG64061 anti-SNAIL antibody at 1:500 dilution.