

ARG64082 anti-CACNB4 antibody

Package: 100 µg
Store at: -20°C

Summary

Product Description	Goat Polyclonal antibody recognizes CACNB4
Tested Reactivity	Hu
Predict Reactivity	Ms, Rat, Dog
Tested Application	WB
Specificity	This antibody is expected to recognise all three reported isoforms (NP_001005747.1; NP_000717.2; NP_001005746.1).
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	CACNB4
Species	Human
Immunogen	C-DYPDSYQDTYKPH
Conjugation	Un-conjugated
Alternate Names	CACNLB4; EJM; EIG9; Calcium channel voltage-dependent subunit beta 4; EJM6; EJM4; Voltage-dependent L-type calcium channel subunit beta-4; EA5; CAB4

Application Instructions

Application table	Application	Dilution
	WB	0.2 - 0.6 µg/ml

Application Note WB: Recommend incubate at RT for 1h.
* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Properties

Form	Liquid
Purification	Purified from goat serum by antigen affinity chromatography.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links

[GeneID: 785 Human](#)

[Swiss-port # O00305 Human](#)

Background

This gene encodes a member of the beta subunit family of voltage-dependent calcium channel complex proteins. Calcium channels mediate the influx of calcium ions into the cell upon membrane polarization and consist of a complex of alpha-1, alpha-2/delta, beta, and gamma subunits in a 1:1:1:1 ratio. Various versions of each of these subunits exist, either expressed from similar genes or the result of alternative splicing. The protein encoded by this locus plays an important role in calcium channel function by modulating G protein inhibition, increasing peak calcium current, controlling the alpha-1 subunit membrane targeting and shifting the voltage dependence of activation and inactivation. Certain mutations in this gene have been associated with idiopathic generalized epilepsy (IGE) and juvenile myoclonic epilepsy (JME). Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar 2009]

Research Area

Signaling Transduction antibody

Calculated Mw

58 kDa

Images



ARG64082 anti-CACNB4 antibody WB image

Western Blot: Human Bone Marrow lysate (35 µg protein in RIPA buffer) stained with ARG64082 anti-CACNB4 antibody at 0.2 µg/ml dilution.