

ARG64120 anti-RBP4 antibody

Package: 100 µg
Store at: -20°C

Summary

Product Description	Goat Polyclonal antibody recognizes RBP4
Tested Reactivity	Hu
Predict Reactivity	Ms, Rat, Dog
Tested Application	WB
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	RBP4
Species	Human
Immunogen	C-DTEDPAKFKMKY
Conjugation	Un-conjugated
Alternate Names	Plasma retinol-binding protein; MCOPCB10; 1-176; RDCCAS; RBP; Retinol-binding protein 4; PRBP; 1-182; 1-179; 1-181

Application Instructions

Application table	Application	Dilution
	WB	0.3 - 1 µg/ml

Application Note
WB: Recommend incubate at RT for 1h.
* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Properties

Form	Liquid
Purification	Purified from goat serum by antigen affinity chromatography.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links

[GeneID: 5950 Human](#)

[Swiss-port # P02753 Human](#)

Background

This protein belongs to the lipocalin family and is the specific carrier for retinol (vitamin A alcohol) in the blood. It delivers retinol from the liver stores to the peripheral tissues. In plasma, the RBP-retinol complex interacts with transthyretin which prevents its loss by filtration through the kidney glomeruli. A deficiency of vitamin A blocks secretion of the binding protein posttranslationally and results in defective delivery and supply to the epidermal cells. [provided by RefSeq, Jul 2008]


Research Area

Cell Biology and Cellular Response antibody; Metabolism antibody; Signaling Transduction antibody

Calculated Mw

23 kDa

Images



ARG64120 anti-RBP4 antibody WB image

Western Blot: Human Liver lysate (35 µg protein in RIPA buffer) stained with ARG64120 anti-RBP4 antibody at 0.3 µg/ml dilution.