

**ARG64146**  
anti-STS2 / TULA antibodyPackage: 100 µg  
Store at: -20°C

### Summary

Product Description	Goat Polyclonal antibody recognizes STS2 / TULA
Tested Reactivity	Hu
Predict Reactivity	Dog
Tested Application	IHC-P
Specificity	This antibody is expected to recognize both reported isoforms (NP_061834.1 and NP_001001895.1).
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	STS2 / TULA
Species	Human
Immunogen	QQCSTPDGKYRDP
Conjugation	Un-conjugated
Alternate Names	TULA; Cbl-interacting protein 4; STS-2; Ubiquitin-associated and SH3 domain-containing protein A; T-cell ubiquitin ligand 1; TULA-1; Suppressor of T-cell receptor signaling 2; CLIP4

### Application Instructions

Application table	Application	Dilution
	IHC-P	2.5 µg/ml
Application Note	IHC-P: Antigen Retrieval: Steam tissue section in Citrate buffer (pH 6.0). * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties

Form	Liquid
Purification	Purified from goat serum by antigen affinity chromatography.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

---

Database links [GeneID: 53347 Human](#)

[Swiss-port # P57075 Human](#)


**Background** This gene encodes one of two family members belonging to the T-cell ubiquitin ligand (TULA) family. Both family members can negatively regulate T-cell signaling. This family member can facilitate growth factor withdrawal-induced apoptosis in T cells, which may occur via its interaction with AIF, an apoptosis-inducing factor. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Aug 2011]

**Research Area** Cell Biology and Cellular Response antibody

**Calculated Mw** 74 kDa

## Images

---



ARG64146 anti-STS2 / TULA antibody IHC-P image

Immunohistochemistry: paraffin embedded Human Spleen. (Steamed antigen retrieval with citrate buffer pH 6) stained with ARG64146 anti-STS2 / TULA antibody at 2.5 µg/ml dilution followed by AP-staining.