

Product datasheet

info@arigobio.com

ARG64170 anti-KCNC3 antibody

Package: 100 μg Store at: -20°C

Summary

Product Description Goat Polyclonal antibody recognizes KCNC3

Tested Reactivity Hu
Tested Application WB
Host Goat

Clonality Polyclonal

Isotype IgG

Target Name KCNC3
Species Human

Immunogen C-KPGPPSFLPDLNAN

Conjugation Un-conjugated

Alternate Names KSHIIID; KV3.3; SCA13; Voltage-gated potassium channel subunit Kv3.3; Potassium voltage-gated

channel subfamily C member 3

Application Instructions

Application table	Application	Dilution
	WB	0.3 - 1 μg/ml
Application Note	WB: Recommend incubate at RT for 1h. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations.	

* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Properties

Form Liquid

Purification Purified from goat serum by antigen affinity chromatography.

Buffer Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.

Preservative 0.02% Sodium azide

Stabilizer 0.5% BSA

Concentration 0.5 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links GeneID: 3748 Human

Swiss-port # Q14003 Human

Background The Shaker gene family of Drosophila encodes components of voltage-gated potassium channels and is

comprised of four subfamilies. Based on sequence similarity, this gene is similar to one of these subfamilies, namely the Shaw subfamily. The protein encoded by this gene belongs to the delayed rectifier class of channel proteins and is an integral membrane protein that mediates the voltage-dependent potassium ion permeability of excitable membranes. [provided by RefSeq, Jul 2008]

Research Area Neuroscience antibody

Calculated Mw 81 kDa

PTM N-glycosylated.

Images

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

ARG64170 anti-KCNC3 antibody WB image

Western Blot: human brain (Frontal Cortex) lysate (35 μg protein in RIPA buffer) stained with ARG64170 anti-KCNC3 antibody at 0.3 $\mu g/ml$ dilution.