

ARG64194 anti-NANOG antibody

Package: 100 µg
Store at: -20°C

Summary

Product Description	Goat Polyclonal antibody recognizes NANOG
Tested Reactivity	Hu
Tested Application	IHC-P, WB
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	NANOG
Species	Human
Immunogen	C-DLFLNYSMNMQPED
Conjugation	Un-conjugated
Alternate Names	Homeobox transcription factor Nanog; Homeobox protein NANOG; hNanog

Application Instructions

Application table	Application	Dilution
	IHC-P	1 - 2 µg/ml
	WB	0.05 - 0.1 µg/ml
Application Note	IHC-P: Antigen Retrieval: Steam tissue section in Citrate buffer (pH 6.0). WB: Recommend incubate at RT for 1h. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification	Purified from goat serum by antigen affinity chromatography.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Database links	GeneID: 79923 Human Swiss-port # Q9H9S0 Human
Gene Symbol	NANOG
Gene Full Name	Nanog homeobox
Background	The protein encoded by this gene is a DNA binding homeobox transcription factor involved in embryonic stem (ES) cell proliferation, renewal, and pluripotency. The encoded protein can block ES cell differentiation and can also autorepress its own expression in differentiating cells. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2015]
Function	Transcription regulator involved in inner cell mass and embryonic stem (ES) cells proliferation and self-renewal. Imposes pluripotency on ES cells and prevents their differentiation towards extraembryonic endoderm and trophoctoderm lineages. Blocks bone morphogenetic protein-induced mesoderm differentiation of ES cells by physically interacting with SMAD1 and interfering with the recruitment of coactivators to the active SMAD transcriptional complexes. Acts as a transcriptional activator or repressor. Binds optimally to the DNA consensus sequence 5'-TAAT[GT][GT]-3' or 5'-[CG][GA][CG]C[GC]ATTAN[GC]-3'. Able to autorepress its expression in differentiating (ES) cells: binds to its own promoter following interaction with ZNF281/ZFP281, leading to recruitment of the NuRD complex and subsequent repression of expression. When overexpressed, promotes cells to enter into S phase and proliferation. [UniProt]
Highlight	Related products: NANOG antibodies ; Anti-Goat IgG secondary antibodies ; Related news: Stem cell and the regenerative medicine: Ready for the patients
Research Area	Cancer antibody; Developmental Biology antibody; Gene Regulation antibody
Calculated Mw	35 kDa

Images



ARG64194 anti-NANOG antibody WB image

Western Blot: HeLa (nuclear) cell lysate (35 µg protein in RIPA buffer) stained with ARG64194 anti-NANOG antibody at 0.05 µg/ml dilution.



ARG64194 anti-NANOG antibody IHC-P image

Immunohistochemistry: paraffin embedded Human Testis. (Steamed antigen retrieval with citrate buffer pH 6) stained with ARG64194 anti-NANOG antibody at 2 µg/ml dilution followed by AP-staining.