

## ARG64254 anti-GAB2 antibody

Package: 100 µg  
Store at: -20°C

### Summary

Product Description	Goat Polyclonal antibody recognizes GAB2
Tested Reactivity	Hu
Predict Reactivity	Ms, Rat, Dog
Tested Application	WB
Specificity	This antibody is expected to recognize both reported isoforms (NP_536739.1; NP_036428.1).
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	GAB2
Species	Human
Immunogen	C-QVDKEKTQALQN
Conjugation	Un-conjugated
Alternate Names	GRB2-associated binder 2; Growth factor receptor bound protein 2-associated protein 2; GRB2-associated-binding protein 2; pp100

### Application Instructions

Application table	Application	Dilution
	WB	0.5 - 1.5 µg/ml
Application Note	WB: Recommend incubate at RT for 1h. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties

Form	Liquid
Purification	Purified from goat serum by antigen affinity chromatography.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note

For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

Database links

[GeneID: 9846 Human](#)

[Swiss-port # Q9UQC2 Human](#)

Background

This gene is a member of the GRB2-associated binding protein (GAB) gene family. These proteins contain pleckstrin homology (PH) domain, and bind SHP2 tyrosine phosphatase and GRB2 adapter protein. They act as adapters for transmitting various signals in response to stimuli through cytokine and growth factor receptors, and T- and B-cell antigen receptors. The protein encoded by this gene is the principal activator of phosphatidylinositol-3 kinase in response to activation of the high affinity IgE receptor. Two alternatively spliced transcripts encoding different isoforms have been described for this gene. [provided by RefSeq, Nov 2009]

Research Area

Cancer antibody; Neuroscience antibody; Signaling Transduction antibody

Calculated Mw

74 kDa

PTM

Phosphorylated on tyrosine residue(s) by the thrombopoietin receptor (TPOR), stem cell factor receptor (SCFR), and T-cell and B-cell antigen receptors, gp130, IL-2R and IL-3R (By similarity). Phosphorylated upon stimulation of TNFRSF11A/RANK by TNFSF11/RANKL (By similarity). Phosphorylated upon EGF stimulation. Phosphorylated on tyrosine residues by HCK upon IL6 signaling. Dephosphorylated by PTPN11.

## Images



ARG64254 anti-GAB2 antibody WB image

Western Blot: K562 cell lysate (35 µg protein in RIPA buffer) stained with ARG64254 anti-GAB2 antibody at 0.5 µg/ml dilution.