

## ARG64385 anti-RPL22 antibody

Package: 100 µg  
Store at: -20°C

### Summary

Product Description	Goat Polyclonal antibody recognizes RPL22
Tested Reactivity	Hu
Predict Reactivity	Ms, Rat
Tested Application	IHC-P
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	RPL22
Species	Human
Immunogen	C-SKESYELRYFQINQ
Conjugation	Un-conjugated
Alternate Names	Epstein-Barr virus small RNA-associated protein; EBER-associated protein; 60S ribosomal protein L22; HBP15; L22; Heparin-binding protein HBp15; EAP; HBP15/L22

### Application Instructions


Application table	Application	Dilution
	IHC-P	3 - 6 µg/ml
Application Note	IHC-P: Antigen Retrieval: Steam tissue section in Citrate buffer (pH 6.0). * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties

Form	Liquid
Purification	Purified from goat serum by antigen affinity chromatography.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.


Database links	<a href="#">GeneID: 6146 Human</a> <a href="#">Swiss-port # P35268 Human</a>
Background	Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a cytoplasmic ribosomal protein that is a component of the 60S subunit. The protein belongs to the L22E family of ribosomal proteins. Its initiating methionine residue is post-translationally removed. The protein can bind specifically to Epstein-Barr virus-encoded RNAs (EBERs) 1 and 2. The mouse protein has been shown to be capable of binding to heparin. Transcript variants utilizing alternative polyA signals exist. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. It was previously thought that this gene mapped to 3q26 and that it was fused to the acute myeloid leukemia 1 (AML1) gene located at 21q22 in some therapy-related myelodysplastic syndrome patients with 3;21 translocations; however, these fusions actually involve a ribosomal protein L22 pseudogene located at 3q26, and this gene actually maps to 1p36.3-p36.2. [provided by RefSeq, Jul 2008]
Research Area	Gene Regulation antibody
Calculated Mw	15 kDa

Images




ARG64385 anti-RPL22 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human thyroid tissue. Antigen Retrieval: Steam tissue section in Citrate buffer (pH 6.0). The tissue section was stained with ARG64385 anti-RPL22 antibody at 3.75 µg/ml dilution followed by AP-staining.




ARG64385 anti-RPL22 antibody IHC-P image

Immunohistochemistry: Paraffin embedded Human Cortex (Steamed antigen retrieval with citrate buffer pH 6) stained with ARG64385 anti-RPL22 antibody at 3.8 µg/ml dilution followed by AP-staining.



ARG64385 anti-RPL22 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human uterus tissue.  
Antigen Retrieval: Steam tissue section in Citrate buffer (pH 6.0). The tissue section was stained with ARG64385 anti-RPL22 antibody at 3.75 µg/ml dilution followed by AP-staining.



ARG64385 anti-RPL22 antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human testis tissue.  
Antigen Retrieval: Steam tissue section in Citrate buffer (pH 6.0). The tissue section was stained with ARG64385 anti-RPL22 antibody at 3.75 µg/ml dilution followed by AP-staining.