

ARG64451 anti-GOT2 antibody

Package: 100 µg
Store at: -20°C

Summary

Product Description	Goat Polyclonal antibody recognizes GOT2
Tested Reactivity	Hu
Predict Reactivity	Ms, Rat
Tested Application	WB
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	GOT2
Species	Human
Immunogen	CKDADEAKRVES
Conjugation	Un-conjugated
Alternate Names	Kynurenine aminotransferase 4; mitAAT; Plasma membrane-associated fatty acid-binding protein; Kynurenine--oxoglutarate transaminase 4; Glutamate oxaloacetate transaminase 2; KAT4; Transaminase A; KATIV; FABP-1; Aspartate aminotransferase, mitochondrial; FABPpm; Fatty acid-binding protein; Kynurenine--oxoglutarate transaminase IV; mAspAT; Kynurenine aminotransferase IV; EC 2.6.1.1; EC 2.6.1.7

Application Instructions

Application table	Application	Dilution
	WB	0.03 - 0.1 µg/ml

Application Note
WB: Recommend incubate at RT for 1h.
* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Properties

Form	Liquid
Purification	Purified from goat serum by antigen affinity chromatography.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links

[GeneID: 2806 Human](#)

[Swiss-port # P00505 Human](#)

Background

Glutamic-oxaloacetic transaminase is a pyridoxal phosphate-dependent enzyme which exists in cytoplasmic and inner-membrane mitochondrial forms, GOT1 and GOT2, respectively. GOT plays a role in amino acid metabolism and the urea and tricarboxylic acid cycles. The two enzymes are homodimeric and show close homology. [provided by RefSeq, Jul 2008]

Research Area

Cancer antibody; Metabolism antibody; Signaling Transduction antibody

Calculated Mw

48 kDa

Images



ARG64451 anti-GOT2 antibody WB image

Western Blot: Human Kidney lysate (35 µg protein in RIPA buffer) stained with ARG64451 anti-GOT2 (aa 295 to 306) antibody at 0.03 µg/ml dilution.