

ARG64479
anti-DACH2 / Dachshund homolog 2 antibodyPackage: 100 µg
Store at: -20°C

Summary

Product Description	Goat Polyclonal antibody recognizes DACH2 / Dachshund homolog 2
Tested Reactivity	Ms
Predict Reactivity	Hu
Tested Application	WB
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	DACH2 / Dachshund homolog 2
Species	Human
Immunogen	C-TRKQAVNSSRPGR
Conjugation	Un-conjugated
Alternate Names	Dach2; Dachshund homolog 2

Application Instructions

Application table	Application	Dilution
	WB	0.03 - 0.1 µg/ml

Application Note WB: Recommend incubate at RT for 1h.
* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Properties

Form	Liquid
Purification	Purified from goat serum by antigen affinity chromatography.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links [GeneID: 93837 Mouse](#)

[Swiss-port # Q925Q8 Mouse](#)

Background

This gene is one of two genes which encode a protein similar to the Drosophila protein dachshund, a transcription factor involved in cell fate determination in the eye, limb and genital disc of the fly. The encoded protein contains two characteristic dachshund domains: an N-terminal domain responsible for DNA binding and a C-terminal domain responsible for protein-protein interactions. This gene is located on the X chromosome and is subject to inactivation by DNA methylation. The encoded protein may be involved in regulation of organogenesis and myogenesis, and may play a role in premature ovarian failure. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Nov 2008]

Research Area Gene Regulation antibody

Calculated Mw 65 kDa

Images



ARG64479 anti-DACH2 / Dachshund homolog 2 antibody WB image

Western blot: NIH3T3 cell lysate (35 µg protein in RIPA buffer) stained with ARG64479 anti-DACH2 / Dachshund homolog 2 antibody at 0.03 µg/ml dilution.