

## ARG64942 anti-NOXA antibody

Package: 100 µg, 50 µg  
Store at: -20°C

### Summary

Product Description	Goat Polyclonal antibody recognizes NOXA
Tested Reactivity	Hu
Tested Application	WB
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	NOXA
Species	Human
Immunogen	C-TQLRRFGDKLNFRQK
Conjugation	Un-conjugated
Alternate Names	NOXA; APR; Phorbol-12-myristate-13-acetate-induced protein 1; PMA-induced protein 1; Immediate-early-response protein APR; Protein Noxa

### Application Instructions

Application table	Application	Dilution
	WB	0.03 - 0.1 µg/ml
Application Note	WB: Recommend incubate at RT for 1h. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties

Form	Liquid
Purification	Purified from goat serum by antigen affinity chromatography.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

---

Database links	<a href="#">GeneID: 5366 Human</a> <a href="#">Swiss-port # Q13794 Human</a>
Gene Symbol	PMAIP1
Gene Full Name	phorbol-12-myristate-13-acetate-induced protein 1
Function	Promotes activation of caspases and apoptosis. Promotes mitochondrial membrane changes and efflux of apoptogenic proteins from the mitochondria. Contributes to p53/TP53-dependent apoptosis after radiation exposure. Promotes proteasomal degradation of MCL1. Competes with BAK1 for binding to MCL1 and can displace BAK1 from its binding site on MCL1 (By similarity). Competes with BIM/BCL2L11 for binding to MCL1 and can displace BIM/BCL2L11 from its binding site on MCL1. [UniProt]
Research Area	Cancer antibody; Cell Biology and Cellular Response antibody; Cell Death antibody; Metabolism antibody
Calculated Mw	6 kDa

## Images

---



ARG64942 anti-NOXA antibody WB image

Western Blot: Human Cerebellum lysate (35 µg protein in RIPA buffer) stained with ARG64942 anti-NOXA antibody at 0.03 µg/ml dilution.