

## Product datasheet

info@arigobio.com

# ARG65023 anti-DSG1 / Desmoglein 1 antibody

Package: 100 μg Store at: -20°C

#### **Summary**

Product Description Goat Polyclonal antibody recognizes DSG1 / Desmoglein 1

Tested Reactivity Hu
Tested Application IHC-P
Host Goat

**Clonality** Polyclonal

Isotype IgG

Target Name DSG1 / Desmoglein 1

Species Human

Immunogen C-YTTSDTLKPSVHVHD

Conjugation Un-conjugated

Alternate Names PPKS1; Desmoglein-1; SPPK1; DGI; EPKHIA; CDHF4; DG1; Desmosomal glycoprotein 1; DSG; Cadherin

family member 4; Pemphigus foliaceus antigen; EPKHE

### **Application Instructions**

Application	Dilution	
IHC-P	4 - 6 μg/ml	
* The dilutions indicate	IHC-P: Antigen Retrieval: Steam tissue section in Citrate buffer (pH 6.0).  * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations	
	IHC-P: Antigen Retrieval * The dilutions indicate	IHC-P: Antigen Retrieval: Steam tissue section in Citrate buffer (pH 6.0).

#### **Properties**

Form Liquid

**Purification** Purified from goat serum by antigen affinity chromatography.

Buffer Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.

Preservative 0.02% Sodium azide

Stabilizer 0.5% BSA

Concentration 0.5 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed

before use.

Note For laboratory research only, not for drug, diagnostic or other use.

#### Bioinformation

Database links GeneID: 1828 Human

Swiss-port # Q02413 Human

Background Desmosomes are cell-cell junctions between epithelial, myocardial and certain other cell types.


Desmoglein 1 is a calcium-binding transmembrane glycoprotein component of desmosomes in vertebrate epithelial cells. Currently, three desmoglein subfamily members have been identified and all are members of the cadherin cell adhesion molecule superfamily. These desmoglein gene family members are located in a cluster on chromosome 18. The protein encoded by this gene has been identified as the autoantigen of the autoimmune skin blistering disease pemphigus foliaceus. [provided

by RefSeq, Jul 2008]

Research Area Signaling Transduction antibody

Calculated Mw 114 kDa

#### **Images**



#### ARG65023 anti-DSG1 / Desmoglein 1 antibody IHC-P image

Immunohistochemistry: Paraffin embedded Human Skin. (Steamed antigen retrieval with citrate buffer pH 6) stained with ARG65023 anti-DSG1 / Desmoglein 1 antibody at 3.8  $\mu$ g/ml dilution followed by AP-staining.