

ARG65322 anti-ZAP70 antibody

Package: 50 µg, 25 µg
Store at: -20°C

Summary

Product Description	Goat Polyclonal antibody recognizes ZAP70
Tested Reactivity	Hu
Predict Reactivity	Ms, Rat, Cow, Dog
Tested Application	IP, WB
Specificity	This antibody is expected to recognize isoform 1 (NP_001070.2) only.
Host	Goat
Clonality	Polyclonal
Isotype	IgG
Target Name	ZAP70
Species	Human
Immunogen	Peptide with sequence C-SDGYTPEPARITSPD, from the internal region of the protein sequence according to NP_001070.2.
Conjugation	Un-conjugated
Alternate Names	STD; SRK; STCD; 70 kDa zeta-chain associated protein; Tyrosine-protein kinase ZAP-70; TZK; Syk-related tyrosine kinase; ZAP-70; EC 2.7.10.2

Application Instructions

Application table	Application	Dilution
	IP	Assay - dependent
	WB	0.3 - 1 µg/ml

Application Note
WB: Recommend incubate at RT for 1h.
* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

Properties


Form	Liquid
Purification	Purified from goat serum by antigen affinity chromatography.
Buffer	Tris saline (pH 7.3), 0.02% Sodium azide and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	0.5% BSA
Concentration	0.5 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated

freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links	GeneID: 7535 Human Swiss-port # P43403 Human
Gene Symbol	ZAP70
Gene Full Name	zeta chain of T cell receptor associated protein kinase 70
Background	This gene encodes an enzyme belonging to the protein tyrosine kinase family, and it plays a role in T-cell development and lymphocyte activation. This enzyme, which is phosphorylated on tyrosine residues upon T-cell antigen receptor (TCR) stimulation, functions in the initial step of TCR-mediated signal transduction in combination with the Src family kinases, Lck and Fyn. This enzyme is also essential for thymocyte development. Mutations in this gene cause selective T-cell defect, a severe combined immunodeficiency disease characterized by a selective absence of CD8-positive T-cells. Two transcript variants that encode different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]
Function	Tyrosine kinase that plays an essential role in regulation of the adaptive immune response. Regulates motility, adhesion and cytokine expression of mature T-cells, as well as thymocyte development. Contributes also to the development and activation of primary B-lymphocytes. When antigen presenting cells (APC) activate T-cell receptor (TCR), a serie of phosphorylations lead to the recruitment of ZAP70 to the doubly phosphorylated TCR component CD247/CD3Z through ITAM motif at the plasma membrane. This recruitment serves to localization to the stimulated TCR and to relieve its autoinhibited conformation. Release of ZAP70 active conformation is further stabilized by phosphorylation mediated by LCK. Subsequently, ZAP70 phosphorylates at least 2 essential adapter proteins: LAT and LCP2. In turn, a large number of signaling molecules are recruited and ultimately lead to lymphokine production, T-cell proliferation and differentiation. Furthermore, ZAP70 controls cytoskeleton modifications, adhesion and mobility of T-lymphocytes, thus ensuring correct delivery of effectors to the APC. ZAP70 is also required for TCR-CD247/CD3Z internalization and degradation through interaction with the E3 ubiquitin-protein ligase CBL and adapter proteins SLA and SLA2. Thus, ZAP70 regulates both T-cell activation switch on and switch off by modulating TCR expression at the T-cell surface. During thymocyte development, ZAP70 promotes survival and cell-cycle progression of developing thymocytes before positive selection (when cells are still CD4/CD8 double negative). Additionally, ZAP70-dependent signaling pathway may also contribute to primary B-cells formation and activation through B-cell receptor (BCR). [UniProt]
Highlight	Related Antibody Duos and Panels: ARG30033 Phospho ZAP70 Antibody Duo (Total, pY319) ARG30049 SyK / Zap70 Antibody Duo Related products: ZAP70 antibodies; ZAP70 Duos / Panels; Anti-Goat IgG secondary antibodies;
Research Area	Controls and Markers antibody; Immune System antibody; Signaling Transduction antibody; SyK / Zap70 Pathway antibody
Calculated Mw	70 kDa
PTM	Phosphorylated on tyrosine residues upon T-cell antigen receptor (TCR) stimulation. Phosphorylation of Tyr-315 and Tyr-319 are essential for ZAP70 positive function on T-lymphocyte activation whereas Tyr-292 has a negative regulatory role. Within the C-terminal kinase domain, Tyr-492 and Tyr-493 are phosphorylated after TCR induction, Tyr-492 playing a negative regulatory role and Tyr-493 a positive. Tyr-493 is dephosphorylated by PTN22. Ubiquitinated in response to T cell activation. Deubiquitinated by OTUD7B.



ARG65322 anti-ZAP70 antibody WB image

Western blot: 30 µg of Jurkat and Molt4 cell lysates stained with ARG65322 anti-ZAP70 antibody at 1:500 dilution.