

Product datasheet

info@arigobio.com

ARG65396 anti-IFN gamma antibody [4S.B3] (Biotin)

Package: 50 μg Store at: 4°C

Summary

Product Description Biotin-conjugated Mouse Monoclonal antibody [4S.B3] recognizes IFN gamma

Tested Reactivity Hu, NHuPrm

Tested Application ELISA, FACS, ICC/IF, IHC-P, WB

Specificity The clone 4S.B3 recognizes IFN-gamma, a 16-25 kDa cytokine produced by activated Th1 cells and NK

cells. Binds both glycosylated and non-glycosylated protein.

Host Mouse

Clonality Monoclonal

Clone 4S.B3

Isotype IgG1

Target Name IFN gamma

Species Human

Immunogen Interferon gamma derived from human leukocytes

Conjugation Biotin

Alternate Names IFN-gamma; Interferon gamma; Immune interferon; IFG; IFI

Application Instructions

Application table	Application	Dilution
	ELISA	0.5 - 2 μg/ml
	FACS	1 - 4 μg/ml
	ICC/IF	Assay-dependent
	IHC-P	Assay-dependent
	WB	Assay-dependent
Application Note	ELISA: ELISA is being used as detection antibody in combination with capture antibody clone NIB42 Flow Cytometry: permeabilization step is suggested before staining. * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form	Liquid
Purification Note	The purified antibody is conjugated with Biotin-LC-NHS under optimum conditions. The reagent is free of unconjugated biotin.
Buffer	PBS (pH 7.4), 15 mM Sodium azide and 0.2% BSA.

Preservative 15 mM Sodium azide

Stabilizer 0.2% BSA

Concentration 1 mg/ml

Storage instruction Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid

repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be

gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Database links GeneID: 3458 Human

Swiss-port # P01579 Human

Background The Interferon gamma (IFN-gamma; 16-25 kDa) is an important regulator of the immune response,

produced in activated Th1 cells and NK cells, particularly in response to IL-2, TNF-alpha and IL-12; its production is suppressed by IL-4, IL-10, and TGF-beta. The producing of IFN-gamma is activated by specific antigens or mitogens through the T cell antigen receptor. IFN-gamma polypeptide forms: 40-60 kDa forms are observable under non-denaturing conditions as dimers and trimers; 20 kDa and 25 kDa forms exist due to variable glycosylation. IFN-gamma belongs to the type II interferons, also called

immune IFN.

IFN-gamma shows antiviral activity and has important immunoregulatory functions. It is a potent activator of macrophages and had antiproliferative effects on transformed cells. IFN-gamma plays an

important role in regulating B cell differentiation by simultaneously stimulating class switch

recombination to the IgG3 and IgG2a isotypes while represing class switch recombination to the IgE and IgG1 isotypes. It also appears to promote antigen presentation by B cells through its effects on MHC. Binding of IFN-gamma to its receptor increases the expression of class I MHC on all somatic cells. It also enhances the expression of class II MHC on antigen-presenting cells. IFN-gamma is the major means by which T cells activate macrophages, increasing their ability to kill bacteria, parasites, and tumours. The activation of macrophages by IFN-gamma is essential for the elimination of bacteria that replicate within the phagosomes of macrophages (f.e. Mycobacteria and Listeria monocytogenes). IFN-gamma can potentiate the high antiviral and antitumor effects of the type I interferons (IFN-alpha, IFN-beta).

IFN-gamma may also activate neutrophils and NK cells.

Function Produced by lymphocytes activated by specific antigens or mitogens. IFN-gamma, in addition to having

antiviral activity, has important immunoregulatory functions. It is a potent activator of macrophages, it has antiproliferative effects on transformed cells and it can potentiate the antiviral and antitumor

effects of the type I interferons. [UniProt]

Highlight Related Antibody Duos and Panels:

ARG30234 IFN gamma ELISA Antibody Duo

Related products:

IFN gamma antibodies; IFN gamma ELISA Kits; IFN gamma Duos / Panels; IFN gamma recombinant

proteins; Anti-Mouse IgG secondary antibodies;

Related news:

HMGB1 in inflammation Inflammatory Cytokines


Research Area Cancer antibody; Developmental Biology antibody; Immune System antibody; Signaling Transduction

antibody

Calculated Mw 19 kDa

PTM Proteolytic processing produces C-terminal heterogeneity, with proteins ending alternatively at

Gly-150, Met-157 or Gly-161.



ARG65396 anti-IFN gamma antibody [4S.B3] (Biotin) FACS image

Flow Cytometry: Human peripheral whole blood stained with ARG65396 anti-IFN gamma antibody [4S.B3] (Biotin).