

**ARG65492**  
**anti-CD105 / Endoglin antibody [MEM-229] (Biotin)**Package: 50 µg  
Store at: 4°C

### Summary

Product Description	Biotin-conjugated Mouse Monoclonal antibody [MEM-229] recognizes CD105 / Endoglin
Tested Reactivity	Hu, Pig
Tested Application	FACS
Specificity	The clone MEM-229 recognizes CD105 (Endoglin), a 90 kDa type I integral membrane homodimer glycoprotein expressed on vascular endothelial cells (small and large vessels), activated monocytes and tissue macrophages, stromal cells of certain tissues including bone marrow, pre-B lymphocytes in fetal marrow and erythroid precursors in fetal and adult bone marrow; it is also present on syncytiotrophoblast on placenta throughout pregnancy.
Host	Mouse
Clonality	Monoclonal
Clone	MEM-229
Isotype	IgG2a
Target Name	CD105 / Endoglin
Species	Human
Immunogen	Recombinant Vaccinia virus containing the human CD105 (L-isoform) cDNA.
Conjugation	Biotin
Alternate Names	CD antigen CD105; HHT1; Endoglin; ORW1; END

### Application Instructions

Application table	Application	Dilution
	FACS	8 µg/ml

**Application Note** \* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.

### Properties

Form	Liquid
Purification Note	The purified antibody is conjugated with Biotin-LC-NHS under optimum conditions. The reagent is free of unconjugated biotin.
Buffer	PBS (pH 7.4), 15 mM Sodium azide and 0.2% BSA.
Preservative	15 mM Sodium azide
Stabilizer	0.2% BSA
Concentration	1 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid


repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

Database links	<a href="#">GeneID: 2022 Human</a> <a href="#">GeneID: 397096 Pig</a> <a href="#">Swiss-port # P17813 Human</a> <a href="#">Swiss-port # P37176 Pig</a>
Gene Symbol	ENG
Gene Full Name	endoglin
Background	CD105 (Endoglin) is a homodimeric transmembrane glycoprotein serving in presence of TGFbetaR-2 as a receptor for TGFbeta-1 and TGFbeta-3. CD105 is highly expressed on endothelial cells and promotes angiogenesis during wound healing, infarcts and in a wide range of tumours and its gene expression is stimulated by hypoxia. CD105 prevents apoptosis in hypoxic endothelial cells and also antagonises the inhibitory effects of TGFbeta-1 on vascular endothelial cell growth and migration. Normal cellular levels of CD105 are required for formation of new blood vessels.
Function	Major glycoprotein of vascular endothelium. Involved in the regulation of angiogenesis. May play a critical role in the binding of endothelial cells to integrins and/or other RGD receptors. Acts as TGF-beta coreceptor and is involved in the TGF-beta/BMP signaling cascade. Required for GDF2/BMP9 signaling through SMAD1 in endothelial cells and modulates TGF-beta1 signaling through SMAD3. [UniProt]
Research Area	Cancer antibody; Cell Biology and Cellular Response antibody; Controls and Markers antibody; Developmental Biology antibody; Immune System antibody
Calculated Mw	71 kDa

## Images



ARG65492 anti-CD105 / Endoglin antibody [MEM-229] (Biotin) FACS image

Flow Cytometry: HUVEC cells (red) and HEK293 cells (blue, negative control) stained with ARG65492 anti-CD105 / Endoglin antibody [MEM-229] (Biotin), followed by Streptavidin (PE).