

ARG65757 anti-Hsp 27 antibody

Package: 100 µl, 50 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes Hsp 27
Tested Reactivity	Hu, Ms, Rat
Tested Application	ICC/IF, IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	Hsp 27
Species	Human
Immunogen	Synthetic peptide derived from Human Hsp 27.
Conjugation	Un-conjugated
Alternate Names	HSP 27; Heat shock 27 kDa protein; HMN2B; HS.76067; SRP27; HEL-S-102; HspB1; CMT2F; 28 kDa heat shock protein; HSP27; Heat shock protein beta-1; Hsp25; Estrogen-regulated 24 kDa protein; Stress-responsive protein 27; HSP28

Application Instructions

Application table	Application	Dilution
	ICC/IF	1:50 - 1:200
	IHC-P	1:50 - 1:200
	WB	1:500 - 1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	HeLa, MCF-7, Mouse lung and Rat lung	
Observed Size	~ 25 kDa	

Properties

Form	Liquid
Purification	Affinity purified.
Buffer	PBS (pH 7.4), 150 mM NaCl, 0.02% Sodium azide and 50% Glycerol.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.


Note

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Gene Symbol	HSPB1
Gene Full Name	heat shock 27kDa protein 1
Background	The protein encoded by this gene is induced by environmental stress and developmental changes. The encoded protein is involved in stress resistance and actin organization and translocates from the cytoplasm to the nucleus upon stress induction. Defects in this gene are a cause of Charcot-Marie-Tooth disease type 2F (CMT2F) and distal hereditary motor neuropathy (dHMN). [provided by RefSeq, Oct 2008]
Function	Involved in stress resistance and actin organization. [UniProt]
Calculated Mw	23 kDa
PTM	Phosphorylated upon exposure to protein kinase C activators and heat shock (PubMed:8325890). Phosphorylation by MAPKAPK2 and MAPKAPK3 in response to stress dissociates HSPB1 from large small heat-shock protein (sHsps) oligomers and impairs its chaperone activity and ability to protect against oxidative stress effectively. Phosphorylation by MAPKAPK5 in response to PKA stimulation induces F-actin rearrangement (PubMed:1332886, PubMed:8093612, PubMed:19166925).

Images




ARG65757 anti-Hsp 27 antibody WB image

Western blot: 30 µg of Mouse lung lysate stained with ARG65757 anti-Hsp 27 antibody at 1:500 dilution.




ARG65757 anti-Hsp 27 antibody WB image

Western blot: 30 µg of Rat lung lysates stained with ARG65757 anti-Hsp 27 antibody at 1:500 dilution.



ARG65757 anti-Hsp 27 antibody WB image

Western blot: HeLa cell lysate stained with ARG65757 anti-Hsp 27 antibody.



ARG65757 anti-Hsp 27 antibody WB image

Western blot: 30 µg of MCF-7 cell lysate stained with ARG65757 anti-Hsp 27 antibody at 1:500 dilution.