

ARG65766 anti-CYP3A4 antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes CYP3A4
Tested Reactivity	Hu, Ms
Tested Application	IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	CYP3A4
Species	Human
Immunogen	Fusion protein of Human CYP3A4
Conjugation	Un-conjugated
Alternate Names	EC 1.14.13.-; Nifedipine oxidase; P450PCN1; EC 1.14.13.32; Taurochenodeoxycholate 6-alpha-hydroxylase; Albendazole sulfoxidase; EC 1.14.13.97; Cytochrome P450 NF-25; CYP11A4; Cytochrome P450 HLP; CYP11A3; EC 1.14.13.157; Cytochrome P450 3A4; Cytochrome P450 3A3; P450C3; EC 1.14.13.67; Quinine 3-monooxygenase; Albendazole monooxygenase; CP34; CYP3A3; CP33; 1,8-cineole 2-exo-monooxygenase; NF-25; HLP; Cytochrome P450-PCN1; CYP3A

Application Instructions

Application table	Application	Dilution
	IHC-P	1:50 - 1:200
	WB	1:200 - 1:1000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Human liver cancer, Mouse liver tissue	

Properties


Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS (pH 7.4), 0.05% Sodium azide and 40% Glycerol
Preservative	0.05% Sodium azide
Stabilizer	40% Glycerol
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Database links	GeneID: 1576 Human Swiss-port # P08684 Human
Gene Symbol	CYP3A4
Gene Full Name	cytochrome P450, family 3, subfamily A, polypeptide 4
Background	This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases that catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by glucocorticoids and some pharmacological agents. This enzyme is involved in the metabolism of approximately half the drugs in use today, including acetaminophen, codeine, cyclosporin A, diazepam and erythromycin. The enzyme also metabolizes some steroids and carcinogens. This gene is part of a cluster of cytochrome P450 genes on chromosome 7q21.1. Previously another CYP3A gene, CYP3A3, was thought to exist; however, it is now thought that this sequence represents a transcript variant of CYP3A4. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Feb 2011]
Function	Cytochromes P450 are a group of heme-thiolate monooxygenases. In liver microsomes, this enzyme is involved in an NADPH-dependent electron transport pathway. It performs a variety of oxidation reactions (e.g. caffeine 8-oxidation, omeprazole sulphoxidation, midazolam 1'-hydroxylation and midazolam 4-hydroxylation) of structurally unrelated compounds, including steroids, fatty acids, and xenobiotics. Acts as a 1,8-cineole 2-exo-monooxygenase. The enzyme also hydroxylates etoposide. [UniProt]
Calculated Mw	57 kDa
PTM	Polyubiquitinated in the presence of AMFR and UBE2G1 and also STUB1/CHIP and UBE2D1 (in vitro).

Images



ARG65766 anti-CYP3A4 antibody IHC-P image

Immunohistochemistry: paraffin-embedded Human liver cancer tissue stained with ARG65766 anti-CYP3A4 antibody (left) at 1:15 dilution, or the same antibody pre-incubated with antigen (right). (Original magnification: X200)



ARG65766 anti-CYP3A4 antibody WB image

Western blot: 40 µg of Mouse liver lysate stained with ARG65766 anti-CYP3A4 antibody at 1:300 dilution.