

## ARG66098 anti-IGFBP3 antibody

Package: 50 µg  
Store at: -20°C

### Summary

Product Description	Rabbit Polyclonal antibody recognizes IGFBP3
Tested Reactivity	Hu
Tested Application	ELISA
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	IGFBP3
Species	Human
Immunogen	E. coli derived recombinant Human IGFBP3. (GASSGGLGPV VRCEPCDARA LAQCAPPNAV CAELVREPGC GCCLTCALSE GQPCGIYTER CGSGLRCQPS PDEARPLQAL LDGRGLCVNA SAVSRLRAYL LPAPPAGNA SESEEDRSAG EVESPSVSST HRVSDPKFHP LHSKIILIKK GHAKDSQRYK VDYESQSTDT QNFSSESKRE TEYGPCRREM EDTLNHLKFL NVLSPRGVHI PNCDDKGFYK KKQCRPSKGR KRGFCWCVDK YGQPLPGYTT KGKEDVHCYS MQSK)
Conjugation	Un-conjugated
Alternate Names	IBP-3; IBP3; Insulin-like growth factor-binding protein 3; IGFBP-3; IGF-binding protein 3; BP-53

### Application Instructions

Application table	Application	Dilution
	ELISA	Sandwich: 0.5 - 2.0 µg/ml with ARG66099 as a detection antibody
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


### Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS (pH 7.2)
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

Database links	<a href="#">GeneID: 3486 Human</a> <a href="#">Swiss-port # P17936 Human</a>
Gene Symbol	IGFBP3
Gene Full Name	insulin-like growth factor binding protein 3
Background	This gene is a member of the insulin-like growth factor binding protein (IGFBP) family and encodes a protein with an IGFBP domain and a thyroglobulin type-I domain. The protein forms a ternary complex with insulin-like growth factor acid-labile subunit (IGFALS) and either insulin-like growth factor (IGF) I or II. In this form, it circulates in the plasma, prolonging the half-life of IGFs and altering their interaction with cell surface receptors. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq, Jul 2008]
Function	IGF-binding proteins prolong the half-life of the IGFs and have been shown to either inhibit or stimulate the growth promoting effects of the IGFs on cell culture. They alter the interaction of IGFs with their cell surface receptors. Also exhibits IGF-independent antiproliferative and apoptotic effects mediated by its receptor TMEM219/IGFBP-3R. [UniProt]
Calculated Mw	32 kDa
PTM	Phosphorylated by FAM20C in the extracellular medium.

## Images



ARG66098 anti-IGFBP3 antibody standard curve image

Sandwich ELISA: ARG66098 anti-IGFBP3 antibody as a capture antibody at 0.5 - 2.0 µg/ml combined with ARG66099 anti-IGFBP3 antibody (Biotin) as a detection antibody. Results of a typical standard run with optical density reading at 405 - 650 nm.