

## ARG66135 anti-CD54 / ICAM1 antibody (Biotin)

Package: 50 µg  
Store at: 4°C

### Summary

Product Description	Biotin-conjugated Rabbit Polyclonal antibody recognizes CD54 / ICAM1
Tested Reactivity	Hu
Tested Application	ELISA, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	CD54 / ICAM1
Species	Human
Immunogen	CHO cells derived recombinant Human CD54 / ICAM1. (QTSVSPSKVI LPRGGSVLVT CSTSCDQPKL LGIETPLPKK ELLPGNNRK VYELSNVQED SQPMCYSNCP DGQSTAKTFL TVYWTPERVE LAPLSWQPV GKNLTLRCQV EGGAPRANLT VVLLRGEKEL KREPAVGPEA EVTITVLVRR DHHGAFSCR TELDLRPQGL ELFENTSAPY QLQTFVLPAT PPQLVSPRVL EVDQTQGTVC SLDGLFPVSE AQVHLALGDQ RLNPVTYGN DSFSKASVS VTADEGTQR LTCAVILGNQ SQETLQVTI YSFPAPNVIL TKPEVSEGE VTKCEAHP AKVTLNGVPA QPLGPRAQLL LKATPEDNGR SFSCSATLEV AGQLIHKNQT RELRVLYGPR LDERDCPGNW TWPENSQQTP MCQAWGNPLP ELKCLKDGT PLPIGESVT TRDLEGTLYC RARSTQGEVT RKVTNVNLS RYE)
Conjugation	Biotin
Alternate Names	CD54; CD antigen CD54; BB2; P3.58; Intercellular adhesion molecule 1; Major group rhinovirus receptor; ICAM-1

### Application Instructions

Application table	Application	Dilution
	ELISA	Direct: 0.25 - 1.0 µg/ml Sandwich: 0.25 - 1.0 µg/ml with ARG66134 as a capture antibody
	WB	0.1 - 0.2 µg/ml
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

### Properties

Form	Liquid
Purification	Purified by affinity chromatography.
Buffer	PBS (pH 7.2)
Concentration	1 mg/ml
Storage instruction	Aliquot and store in the dark at 2-8°C. Keep protected from prolonged exposure to light. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be

gently mixed before use.


## Note

For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation


Database links	<a href="#">GeneID: 3383 Human</a> <a href="#">Swiss-port # P05362 Human</a>
Gene Symbol	ICAM1
Gene Full Name	intercellular adhesion molecule 1
Background	This gene encodes a cell surface glycoprotein which is typically expressed on endothelial cells and cells of the immune system. It binds to integrins of type CD11a / CD18, or CD11b / CD18 and is also exploited by Rhinovirus as a receptor. [provided by RefSeq, Jul 2008]
Function	ICAM proteins are ligands for the leukocyte adhesion protein LFA-1 (integrin alpha-L/beta-2). During leukocyte trans-endothelial migration, ICAM1 engagement promotes the assembly of endothelial apical cups through ARHGEF26/SGEF and RHOG activation (By similarity). [UniProt]
Research Area	Cell Biology and Cellular Response antibody; Developmental Biology antibody; Immune System antibody; Cytotoxic T Cell Surface Study antibody
Calculated Mw	58 kDa
PTM	Monoubiquitinated, which is promoted by MARCH9 and leads to endocytosis.

## Images



ARG66135 anti-CD54 / ICAM1 antibody (Biotin) standard curve image

Direct ELISA: ARG66135 anti-CD54 / ICAM1 antibody (Biotin) at 0.25 - 1.0  $\mu$ g/ml results of a typical standard run with optical density reading at 405 - 650 nm.



ARG66135 anti-CD54 / ICAM1 antibody (Biotin) standard curve image

Sandwich ELISA: ARG66135 anti-CD54 / ICAM1 antibody (Biotin) as a detection antibody at 0.25 - 1.0  $\mu$ g/ml combined with ARG66134 anti-CD54 / ICAM1 antibody as a capture antibody. Results of a typical standard run with optical density reading at 405 - 650 nm.