

## ARG66429 anti-IL4 antibody [SQab18131]

Package: 100 µg  
Store at: -20°C

### Summary

Product Description	Mouse Monoclonal antibody [SQab18131] recognizes IL4
Tested Reactivity	Hu
Tested Application	ELISA, FACS, ICC/IF, IHC-P, WB
Host	Mouse
Clonality	Monoclonal
Clone	SQab18131
Isotype	IgG1, kappa
Target Name	IL4
Species	Human
Immunogen	Recombinant Human IL4 protein.
Conjugation	Un-conjugated
Alternate Names	BSF-1; IL-4; B-cell stimulatory factor 1; BCGF1; Lymphocyte stimulatory factor 1; BCGF-1; Interleukin-4; Pitrakinra; Binetrakin; BSF1

### Application Instructions

Application table	Application	Dilution
	ELISA	1:3000 - 1:10000
	FACS	1:400 - 1:1000
	ICC/IF	1:400 - 1:1000
	IHC-P	Assay-dependent
	WB	1:2500 - 1:5000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	



### Properties

Purification	Affinity purification with immunogen.
Buffer	PBS (pH 7.4) and 0.01% Thimerosal.
Preservative	0.01% Thimerosal
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C or below. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

## Bioinformation


Gene Symbol	IL4
Gene Full Name	interleukin 4
Background	The protein encoded by this gene is a pleiotropic cytokine produced by activated T cells. This cytokine is a ligand for interleukin 4 receptor. The interleukin 4 receptor also binds to IL13, which may contribute to many overlapping functions of this cytokine and IL13. STAT6, a signal transducer and activator of transcription, has been shown to play a central role in mediating the immune regulatory signal of this cytokine. This gene, IL3, IL5, IL13, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL13. This gene, IL13 and IL5 are found to be regulated coordinately by several long-range regulatory elements in an over 120 kilobase range on the chromosome. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]
Function	Participates in at least several B-cell activation processes as well as of other cell types. It is a costimulator of DNA-synthesis. It induces the expression of class II MHC molecules on resting B-cells. It enhances both secretion and cell surface expression of IgE and IgG1. It also regulates the expression of the low affinity Fc receptor for IgE (CD23) on both lymphocytes and monocytes. [UniProt]
Calculated Mw	17 kDa
Cellular Localization	Secreted. [UniProt]

## Images



### ARG66429 anti-IL4 antibody [SQab18131] WB image

Western blot: 100 ng of Recombinant IL4 protein and 60 µg of 293T cell lysate stained with ARG66429 anti-IL4 antibody [SQab18131] at 1:2000 dilution.



### ARG66429 anti-IL4 antibody [SQab18131] ELISA image

ELISA: Titration curve of ARG66429 anti-IL4 antibody [SQab18131]. Red: IL4; Black: BSA (negative control).

