

ARG66562 anti-OPCML / OBCAM antibody

Package: 100 µg
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes OPCML / OBCAM
Tested Reactivity	Mk
Tested Application	WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	OPCML / OBCAM
Species	Human
Immunogen	Synthetic peptide derived from the Internal region of Human OBCAM.
Conjugation	Un-conjugated
Alternate Names	IGLON family member 1; Opioid-binding cell adhesion molecule; IGLON1; OBCAM; OPCM; Opioid-binding protein/cell adhesion molecule; OPCML

Application Instructions

Application table	Application	Dilution
	WB	1:500 - 1:2000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	


Properties

Form	Liquid
Purification	Affinity purification with immunogen.
Buffer	PBS, 0.02% Sodium azide, 50% Glycerol and 0.5% BSA.
Preservative	0.02% Sodium azide
Stabilizer	50% Glycerol and 0.5% BSA
Concentration	1 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	OPCML
Gene Full Name	opioid binding protein/cell adhesion molecule-like
Background	This gene encodes a member of the IgLON subfamily in the immunoglobulin protein superfamily. The encoded protein is localized in the plasma membrane and may have an accessory role in opioid receptor function. This gene has an ortholog in rat and bovine. The opioid binding-cell adhesion molecule encoded by the rat gene binds opioid alkaloids in the presence of acidic lipids, exhibits selectivity for mu ligands and acts as a GPI-anchored protein. Since the encoded protein is highly conserved in species during evolution, it may have a fundamental role in mammalian systems. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]
Function	Binds opioids in the presence of acidic lipids; probably involved in cell contact. [UniProt]
Calculated Mw	38 kDa
Cellular Localization	Cell membrane; Lipid-anchor, GPI-anchor. [UniProt]

Images



ARG66562 anti-OPCML / OBCAM antibody WB image

Western blot: COS7 cell lysate stained with ARG66562 anti-OPCML / OBCAM antibody at 1:1000 dilution.