

Product datasheet

info@arigobio.com

ARG66627 anti-RASGRF1 / CDC25 phospho (Ser916) antibody

Package: 100 μg Store at: -20°C

Summary

Product Description Rabbit Polyclonal antibody recognizes RASGRF1 / CDC25 phospho (Ser916)

Tested Reactivity Hu

Tested Application IHC-P

Host Rabbit

Clonality Polyclonal

Isotype IgG

Target Name RASGRF1 / CDC25

Species Human

Immunogen Phosphospecific peptide around Ser916 of Human RASGRF1 / CDC25.

Conjugation Un-conjugated

Alternate Names CDC25L; Ras-specific guanine nucleotide-releasing factor 1; CDC25; GRF1; PP13187; GRF55; H-GRF55;

Ras-specific nucleotide exchange factor CDC25; Guanine nucleotide-releasing protein; Ras-GRF1; GNRP;

ras-GRF1

Application Instructions

Application table	Application	Dilution
	IHC-P	1:100 - 1:300
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	

Properties

Form Liquid

Purification Affinity purification with immunogen.

Buffer PBS, 0.02% Sodium azide, 50% Glycerol and 0.5% BSA.

Preservative 0.02% Sodium azide

Stabilizer 50% Glycerol and 0.5% BSA

Concentration 1 mg/ml

Storage instruction For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot

and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol RASGRF1

Gene Full Name Ras protein-specific guanine nucleotide-releasing factor 1

Background The protein encoded by this gene is a guanine nucleotide exchange factor (GEF) similar to the

Saccharomyces cerevisiae CDC25 gene product. Functional analysis has demonstrated that this protein stimulates the dissociation of GDP from RAS protein. The studies of the similar gene in mouse

stimulates the dissociation of GDP from RAS protein. The studies of the similar gene in mouse suggested that the Ras-GEF activity of this protein in brain can be activated by Ca2+ influx, muscarinic receptors, and G protein beta-gamma subunit. Mouse studies also indicated that the Ras-GEF signaling pathway mediated by this protein may be important for long-term memory. Alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, Mar 2009]

Function Promotes the exchange of Ras-bound GDP by GTP. [UniProt]


Calculated Mw 145 kDa

PTM Phosphorylated by PLK2, leading to ubiquitination and degradation by the proteasome.

Ubiquitinated and degraded following phosphorylation by PLK2.

Phosphorylated by SRC and LCK. Phosphorylation by LCK increases its capacity to stimulate the GDP/GTP exchange on Ras, whereas its phosphorylation by SRC seems not to have an effect on stimulation activity. [UniProt]

Images



ARG66627 anti-RASGRF1 / CDC25 phospho (Ser916) antibody IHC-P image

Immunohistochemistry: Paraffin-embedded Human brain tissue stained with ARG66627 anti-RASGRF1 / CDC25 phospho (Ser916) antibody. The picture on the right is blocked with the phospho peptide.