

ARG66769 anti-SCGB2A2 / Mammaglobin A antibody [SQab20202]

Package: 100 µl
Store at: -20°C

Summary

Product Description	Recombinant Rabbit Monoclonal antibody [SQab20202] recognizes SCGB2A2 / Mammaglobin A
Tested Reactivity	Hu
Tested Application	IHC-P
Host	Rabbit
Clonality	Monoclonal
Clone	SQab20202
Isotype	IgG
Target Name	SCGB2A2 / Mammaglobin A
Species	Human
Immunogen	Recombinant fragment of Human SCGB2A2 / Mammaglobin A.
Conjugation	Un-conjugated
Alternate Names	UGB2; Secretoglobin family 2A member 2; MGB1; Mammaglobin-A; Mammaglobin-1

Application Instructions

Application table	<table><thead><tr><th>Application</th><th>Dilution</th></tr></thead><tbody><tr><td>IHC-P</td><td>1:100 - 1:200</td></tr></tbody></table>	Application	Dilution	IHC-P	1:100 - 1:200
Application	Dilution				
IHC-P	1:100 - 1:200				
Application Note	IHC-P: Antigen Retrieval: Heat mediation was performed in Tris/EDTA buffer (pH 9.0). * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.				
Positive Control	Breast carcinoma tissue.				


Properties

Form	Liquid
Purification	Purification with Protein A.
Buffer	PBS, 0.01% Sodium azide, 40% Glycerol and 0.05% BSA.
Preservative	0.01% Sodium azide
Stabilizer	40% Glycerol and 0.05% BSA
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	SCGB2A2
Gene Full Name	secretoglobin, family 2A, member 2
Calculated Mw	10 kDa

Images



ARG66769 anti-SCGB2A2 / Mammaglobin A antibody [SQab20202]
IHC-P image

Immunohistochemistry: Formalin/PFA-fixed and paraffin-embedded Human breast carcinoma tissue. Antigen Retrieval: Heat mediation was performed in Tris/EDTA buffer (pH 9.0). The tissue section was stained with ARG66769 anti-SCGB2A2 / Mammaglobin A antibody [SQab20202] at 18°C - 25°C for 30 minutes.