

## ARG67088 anti-AMACR / p504S antibody [SQab30349]

Package: 100 µl  
Store at: -20°C

### Summary

Product Description	Recombinant Mouse Monoclonal antibody [SQab30349] recognizes AMACR / p504S
Tested Reactivity	Hu
Tested Application	IHC-P
Host	Mouse
Clonality	Monoclonal
Clone	SQab30349
Isotype	IgG
Target Name	AMACR / p504S
Species	Human
Immunogen	Synthetic peptide of AMACR / p504S.
Conjugation	Un-conjugated
Alternate Names	AMACR; Alpha-Methylacyl-CoA Racemase; P504S; RACE; 2-Methylacyl-CoA Racemase; EC 5.1.99.4; AMACRD; CBAS4; RM

### Application Instructions

Application table	Application	Dilution
	IHC-P	1:100-1:200
Application Note	IHC-P: Antigen Retrieval: Heat mediation was performed in Tris/EDTA buffer (pH 9.0). * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Positive Control	Prostate cancer.	


### Properties

Form	Liquid
Purification	Purification with Protein A.
Buffer	PBS, 0.01% Sodium azide, 40% Glycerol and 0.05% BSA.
Preservative	0.01% Sodium azide
Stabilizer	40% Glycerol and 0.05% BSA
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

## Bioinformation

Gene Symbol	AMACR
Gene Full Name	Alpha-Methylacyl-CoA Racemase
Background	This gene encodes a racemase. The encoded enzyme interconverts pristanoyl-CoA and C27-bile acylCoAs between their (R)- and (S)-stereoisomers. The conversion to the (S)-stereoisomers is necessary for degradation of these substrates by peroxisomal beta-oxidation. Encoded proteins from this locus localize to both mitochondria and peroxisomes. Mutations in this gene may be associated with adult-onset sensorimotor neuropathy, pigmentary retinopathy, and adrenomyeloneuropathy due to defects in bile acid synthesis. Alternatively spliced transcript variants have been described. Read-through transcription also exists between this gene and the upstream neighboring C1QTNF3 (C1q and tumor necrosis factor related protein 3) gene.
Function	Acts only on coenzyme A thioesters, not on free fatty acids, and accepts as substrates a wide range of alpha-methylacyl-CoAs, including pristanoyl-CoA, trihydroxycoprostanoyl-CoA (an intermediate in bile acid synthesis), and arylpropionic acids like the anti-inflammatory drug ibuprofen (2-(4-isobutylphenyl)propionic acid) but neither 3-methyl-branched nor linear-chain acyl-CoAs.
Calculated Mw	42 kDa
PTM	Acetylation
Cellular Localization	Mitochondrion, Peroxisome

## Images



ARG67088 anti-AMACR / p504S antibody [SQab30349] IHC-P image

Immunohistochemistry: Human Prostate cancer stained with ARG67088 anti-AMACR / p504S antibody [SQab30349].