

ARG67099 anti-Perilipin 1 antibody

Package: 100 µl
Store at: -20°C

Summary

Product Description	Rabbit Polyclonal antibody recognizes Perilipin 1
Tested Reactivity	Hu, Ms, Rat
Tested Application	IHC-P, WB
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Target Name	Perilipin 1
Immunogen	Synthetic peptide
Conjugation	Un-conjugated
Alternate Names	Perilipin-1; PER1; Lipid droplet-associated protein; PLIN; FPLD4

Application Instructions

Application table	Application	Dilution
	IHC-P	1:200 - 1:500
	WB	1:1000 - 1:3000
Application Note	* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist.	
Observed Size	~ 68 kDa	


Properties

Form	Liquid
Purification	Affinity purified.
Buffer	100 mM Tris Glycine (pH 7.0), 0.025% ProClin 300, 20% Glycerol and 1% BSA.
Preservative	0.025% ProClin 300
Stabilizer	20% Glycerol and 1% BSA
Concentration	1.26 mg/ml
Storage instruction	For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation


Gene Symbol	PLIN1
Gene Full Name	perilipin 1
Background	The protein encoded by this gene coats lipid storage droplets in adipocytes, thereby protecting them until they can be broken down by hormone-sensitive lipase. The encoded protein is the major cAMP-dependent protein kinase substrate in adipocytes and, when unphosphorylated, may play a role in the inhibition of lipolysis. Alternatively spliced transcript variants varying in the 5' UTR, but encoding the same protein, have been found for this gene. [provided by RefSeq, Feb 2009]
Function	Modulator of adipocyte lipid metabolism. Coats lipid storage droplets to protect them from breakdown by hormone-sensitive lipase (HSL). Its absence may result in leanness. Plays a role in unilocular lipid droplet formation by activating CIDEC. Their interaction promotes lipid droplet enlargement and directional net neutral lipid transfer. May modulate lipolysis and triglyceride levels. [UniProt]
Calculated Mw	56 kDa
PTM	Major cAMP-dependent protein kinase-substrate in adipocytes, also dephosphorylated by PP1. When phosphorylated, may be maximally sensitive to HSL and when unphosphorylated, may play a role in the inhibition of lipolysis, by acting as a barrier in lipid droplet (By similarity). [UniProt]
Cellular Localization	Endoplasmic reticulum. Lipid droplet. Note=Lipid droplet surface-associated. [UniProt]

Images




ARG67099 anti-Perilipin 1 antibody IHC-P image

Immunohistochemistry: Human cancer stained with ARG67099 anti-Perilipin 1 antibody at 1:100 dilution.




ARG67099 anti-Perilipin 1 antibody WB image

Western blot: Human adipose stained with ARG67099 anti-Perilipin 1 antibody at 1: 1500 dilution.



ARG67099 anti-Perilipin 1 antibody IHC-P image

Immunohistochemistry: Mouse adipose stained with ARG67099 anti-Perilipin 1 antibody at 1:200 dilution.



ARG67099 anti-Perilipin 1 antibody WB image

Western blot: Mouse adipose stained with ARG67099 anti-Perilipin 1 antibody at 1: 1500 dilution.