

# Product datasheet

info@arigobio.com

ARG81270 Human CD62P / P-Selectin ELISA Kit Package: 96 wells Store at: 4°C

## Summary

Product Description ARG81270 Human CD62P / P-Selectin ELISA Kit is an Enzyme Immunoassay kit for the quantification of

Human CD62P / P-Selectin in serum, plasma or cell culture supernatants.

Tested Reactivity Hu

Tested Application ELISA

Target Name CD62P / P-Selectin

Conjugation HRP

Conjugation Note Substrate: TMB and read at 450 nm.

Sensitivity 62 pg/ml

Sample Type Serum, plasma or cell culture supernatants.

Standard Range 125 - 8000 pg/ml

Sample Volume  $100 \ \mu l$ 

Alternate Names PADGEM; CD62; Platelet activation dependent granule-external membrane protein; CD62 antigen-like

family member P; GMP140; PSEL; Granule membrane protein 140; CD62P; CD antigen CD62P; Leukocyte-endothelial cell adhesion molecule 3; GRMP; GMP-140; P-selectin; LECAM3

tedhocyte endothendreen danesion morecule 3, Ghivir, Givir 140, 1 Sciectiff, ELo.

# **Application Instructions**

Assay Time ~ 3.5 hours

#### **Properties**

Form 96 well

Storage instruction Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test

reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual

for detail temperatures of the components.

Note For laboratory research only, not for drug, diagnostic or other use.

#### Bioinformation

Gene Symbol SELP

Gene Full Name selectin P (granule membrane protein 140kDa, antigen CD62)

Background This gene encodes a 140 kDa protein that is stored in the alpha-granules of platelets and Weibel-Palade

bodies of endothelial cells. This protein redistributes to the plasma membrane during platelet activation and degranulation and mediates the interaction of activated endothelial cells or platelets with leukocytes. The membrane protein is a calcium-dependent receptor that binds to sialylated forms of Lewis blood group carbohydrate antigens on neutrophils and monocytes. Alternative splice variants

may occur but are not well documented. [provided by RefSeq, Jul 2008]

Function Ca(2+)-dependent receptor for myeloid cells that binds to carbohydrates on neutrophils and


monocytes. Mediates the interaction of activated endothelial cells or platelets with leukocytes. The ligand recognized is sialyl-Lewis X. Mediates rapid rolling of leukocyte rolling over vascular surfaces

during the initial steps in inflammation through interaction with PSGL1. [UniProt]

Highlight Related products:

CD62 antibodies; CD62 ELISA Kits; New ELISA data calculation tool: Simplify the ELISA analysis by GainData

### **Images**



#### ARG81270 Human CD62P / P-Selectin ELISA Kit standard curve image

ARG81270 Human CD62P / P-Selectin ELISA Kit results of a typical standard run with optical density reading at 450 nm.