

ARG81415 Human MMP10 ELISA Kit

Package: 96 wells Store at: 4°C

Component

Cat. No.	Component Name	Package	Temp
ARG81415-001	Antibody-coated microplate	8 X 12 strips	4°C. Unused strips should be sealed tightly in the air-tight pouch.
ARG81415-002	Standard	2 X 10 ng/vial	4°C
ARG81415-003	Standard/Sample diluent	30 ml (Ready to use)	4°C
ARG81415-004	Antibody conjugate concentrate (100X)	1 vial (100 μl)	4°C
ARG81415-005	Antibody diluent buffer	12 ml (Ready to use)	4°C
ARG81415-006	HRP-Streptavidin concentrate (100X)	1 vial (100 μl)	4°C
ARG81415-007	HRP-Streptavidin diluent buffer	12 ml (Ready to use)	4°C
ARG81415-008	25X Wash buffer	20 ml	4°C
ARG81415-009	TMB substrate	10 ml (Ready to use)	4°C (Protect from light)
ARG81415-010	STOP solution	10 ml (Ready to use)	4°C
ARG81415-011	Plate sealer	4 strips	Room temperature

Summary

Product Description	ARG81415 Human MMP10 ELISA Kit is an Enzyme Immunoassay kit for the quantification of Human MMP10 in serum, plasma (heparin) and cell culture supernatants.
Tested Reactivity	Hu
Tested Application	ELISA
Specificity	There is no detectable cross-reactivity with other relevant proteins.
Target Name	MMP10
Conjugation	HRP
Conjugation Note	Substrate: TMB and read at 450 nm.
Sensitivity	39 pg/ml
Sample Type	Serum, plasma (heparin) and cell culture supernatants.
Standard Range	78 - 5000 pg/ml
Sample Volume	100 μΙ

Alternate Names

Application Instructions


SL-2; STMY2; Transin-2; EC 3.4.24.22; Matrix metalloproteinase-10; MMP-10; Stromelysin-2

Properties

Form	96 well
Storage instruction	Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual for detail temperatures of the components.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	MMP10		
Gene Full Name	matrix metallopeptidase 10		
Background	Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, as well as in disease processes, such as arthritis and metastasis. Most MMP's are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. The enzyme encoded by this gene degrades proteoglycans and fibronectin. The gene is part of a cluster of MMP genes which localize to chromosome 11q22.3. [provided by RefSeq, Jul 2008]		
Function	Can degrade fibronectin, gelatins of type I, III, IV, and V; weakly collagens III, IV, and V. Activates procollagenase. [UniProt]		
Highlight	Related products: <u>MMP10 antibodies; MMP10 ELISA Kits;</u> Related news: <u>Detecting MMPs and their non-ECM substrates</u> New ELISA data calculation tool: <u>Simplify the ELISA analysis by GainData</u>		



ARG81415 Human MMP10 ELISA Kit standard curve image

ARG81415 Human MMP10 ELISA Kit results of a typical standard run with optical density reading at 450 nm.