

Product datasheet

info@arigobio.com

ARG81659 Human CCL22 / MDC ELISA Kit

Package: 96 wells Store at: 4°C

Component

Cat. No.	Component Name	Package	Temp
ARG81659-001	Antibody-coated microplate	8 X 12 strips	4°C. Unused strips should be sealed tightly in the air-tight pouch.
ARG81659-002	Standard	2 X 10 ng/vial	4°C
ARG81659-003	Standard/Sample diluent	30 ml (Ready to use)	4°C
ARG81659-004	Antibody conjugate concentrate (100X)	1 vial (100 μl)	4°C
ARG81659-005	Antibody diluent buffer	12 ml (Ready to use)	4°C
ARG81659-006	HRP-Streptavidin concentrate (100X)	1 vial (100 μl)	4°C
ARG81659-007	HRP-Streptavidin diluent buffer	12 ml (Ready to use)	4°C
ARG81659-008	25X Wash buffer	20 ml	4°C
ARG81659-009	TMB substrate	10 ml (Ready to use)	4°C (Protect from light)
ARG81659-010	STOP solution	10 ml (Ready to use)	4°C
ARG81659-011	Plate sealer	4 strips	Room temperature

Summary

Product Description	ARG81659 Human CCL22 /	MDC ELISA Kit is an Enzyme Immunoassay kit for the quantification of	

Human CCL22 / MDC in serum, plasma (heparin, EDTA) and cell culture supernatants.

Tested Reactivity Hu

Tested Application ELISA

Specificity There is no detectable cross-reactivity with other relevant proteins.

Target Name CCL22 / MDC

Conjugation HRP

Conjugation Note Substrate: TMB and read at 450 nm.

Sensitivity 15.6 pg/ml

Sample Type Serum, plasma (heparin, EDTA) and cell culture supernatants.

Standard Range 31.2 - 2000 pg/ml

Sample Volume $100 \ \mu l$

Precision Intra-Assay CV: 6.2%; Inter-Assay CV: 7.5%

Alternate Names CC chemokine STCP-1; Stimulated T-cell chemotactic protein 1; 3-69; Macrophage-derived chemokine;

MDC; DC/B-CK; SCYA22; Small-inducible cytokine A22; ABCD-1; 7-69; 1-69; A-152E5.1; STCP-1; 5-69; C-C

motif chemokine 22

Application Instructions

Assay Time

~ 5 hours

Properties

Form 96 well

Storage instruction Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test

reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual

for detail temperatures of the components.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol CCL22

Gene Full Name chemokine (C-C motif) ligand 22

Background This antimicrobial gene is one of several Cys-Cys (CC) cytokine genes clustered on the q arm of

chromosome 16. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for monocytes, dendritic cells, natural killer cells and for chronically activated T lymphocytes. It also displays a mild activity for primary activated T

lymphocytes and has no chemoattractant activity for neutrophils, eosinophils and resting T

lymphocytes. The product of this gene binds to chemokine receptor CCR4. This chemokine may play a role in the trafficking of activated T lymphocytes to inflammatory sites and other aspects of activated T

lymphocyte physiology. [provided by RefSeq, Sep 2014]

Function May play a role in the trafficking of activated/effector T-lymphocytes to inflammatory sites and other

aspects of activated T-lymphocyte physiology. Chemotactic for monocytes, dendritic cells and natural killer cells. Mild chemoattractant for primary activated T-lymphocytes and a potent chemoattractant for chronically activated T-lymphocytes but has no chemoattractant activity for neutrophils,

eosinophils, and resting T-lymphocytes. Binds to CCR4. Processed forms MDC(3-69), MDC(5-69) and


MDC(7-69) seem not be active. [UniProt]

Highlight Related products:

CCL22 antibodies; CCL22 ELISA Kits; New ELISA data calculation tool: Simplify the ELISA analysis by GainData

PTM The N-terminal processed forms MDC(3-69), MDC(5-69) and MDC(7-69) are produced by proteolytic

cleavage after secretion from monocyte derived dendrocytes. [UniProt]



ARG81659 Human CCL22 / MDC ELISA Kit standard curve image

ARG81659 Human CCL22 / MDC ELISA Kit results of a typical standard run with optical density reading at 450 nm.