

Product datasheet

info@arigobio.com

ARG81843 Mouse Progranulin ELISA Kit

Package: 96 wells Store at: 4°C

Component

Cat. No.	Component Name	Package	Temp
ARG81843-001	Antibody-coated microplate	8 X 12 strips	4°C. Unused strips should be sealed tightly in the air-tight pouch.
ARG81843-002	Standard	2 X 50 ng/vial	4°C
ARG81843-003	Standard/Sample diluent	30 ml (Ready to use)	4°C
ARG81843-004	Antibody conjugate concentrate (100X)	1 vial (100 μl)	4°C
ARG81843-005	Antibody diluent buffer	12 ml (Ready to use)	4°C
ARG81843-006	HRP-Streptavidin concentrate (100X)	1 vial (100 μl)	4°C
ARG81843-007	HRP-Streptavidin diluent buffer	12 ml (Ready to use)	4°C
ARG81843-008	25X Wash buffer	20 ml	4°C
ARG81843-009	TMB substrate	10 ml (Ready to use)	4°C (Protect from light)
ARG81843-010	STOP solution	10 ml (Ready to use)	4°C
ARG81843-011	Plate sealer	4 strips	Room temperature

Summary

Product Description	ARG81843 Mouse Progranulin ELISA Kit is an Enzyme Immunoassay kit for the quantification of Mouse	

Progranulin in serum, plasma (heparin, EDTA) and cell culture supernatants.

Tested Reactivity Ms

Tested Application ELISA

Specificity There is no detectable cross-reactivity with other relevant proteins.

Target Name Progranulin

Conjugation HRP

Conjugation Note Substrate: TMB and read at 450 nm.

Sensitivity 0.39 ng/ml

Sample Type Serum, plasma (heparin, EDTA) and cell culture supernatants.

Standard Range 0.78 - 50 ng/ml

Sample Volume $100 \ \mu l$

Precision Intra-Assay CV: 5.2%; Inter-Assay CV: 5.7%

Alternate Names PEPI; Glycoprotein 88; CLN11; GEP; Proepithelin; Granulin E; Granulin D; Granulins; PGRN; Glycoprotein

of 88 Kda; Granulin G; Progranulin; PCDGF; Granulin B; Granulin C; Granulin A; GP88; Granulin F

Application Instructions

Assay Time

~ 5 hours

Properties

Form

96 well

Storage instruction

Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual

for detail temperatures of the components.

Note

For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol

GRN

Gene Full Name

granulin

Background

Granulins are a family of secreted, glycosylated peptides that are cleaved from a single precursor protein with 7.5 repeats of a highly conserved 12-cysteine granulin/epithelin motif. The 88 kDa precursor protein, progranulin, is also called proepithelin and PC cell-derived growth factor. Cleavage of the signal peptide produces mature granulin which can be further cleaved into a variety of active, 6 kDa peptides. These smaller cleavage products are named granulin A, granulin B, granulin C, etc. Epithelins 1 and 2 are synonymous with granulins A and B, respectively. Both the peptides and intact granulin protein regulate cell growth. However, different members of the granulin protein family may act as inhibitors, stimulators, or have dual actions on cell growth. Granulin family members are important in normal development, wound healing, and tumorigenesis. [provided by RefSeq, Jul 2008]

Function

Granulins have possible cytokine-like activity. They may play a role in inflammation, wound repair, and

tissue remodeling.

Granulin-4 promotes proliferation of the epithelial cell line A431 in culture while granulin-3 acts as an

antagonist to granulin-4, inhibiting the growth. [UniProt]

Highlight


Related products:

Progranulin antibodies; Progranulin ELISA Kits;

New ELISA data calculation tool: Simplify the ELISA analysis by GainData

PTM

Granulins are disulfide bridged. [UniProt]



ARG81843 Mouse Progranulin ELISA Kit standard curve image

ARG81843 Mouse Progranulin ELISA Kit results of a typical standard run with optical density reading at 450 nm.