

# Product datasheet

info@arigobio.com

### ARG82533 Human SPINK5 ELISA Kit

Package: 96 wells Store at: 4°C

## Component

Cat. No.	Component Name	Package	Temp
ARG82533-001	Antibody-coated microplate	8 X 12 strips	4°C. Unused strips should be sealed tightly in the air-tight pouch.
ARG82533-002	Standard	2 X 10 ng/vial	4°C
ARG82533-003	Standard/Sample diluent	30 ml (Ready to use)	4°C
ARG82533-004	Antibody conjugate concentrate (100X)	1 vial (100 μl)	4°C
ARG82533-005	Antibody diluent buffer	12 ml (Ready to use)	4°C
ARG82533-006	HRP-Streptavidin concentrate (100X)	1 vial (100 μl)	4°C
ARG82533-007	HRP-Streptavidin diluent buffer	12 ml (Ready to use)	4°C
ARG82533-008	25X Wash buffer	20 ml	4°C
ARG82533-009	TMB substrate	10 ml (Ready to use)	4°C (Protect from light)
ARG82533-010	STOP solution	10 ml (Ready to use)	4°C
ARG82533-011	Plate sealer	4 strips	Room temperature

#### Summary

Product Description	ARG82533 Human SPINK5 ELISA Kit is an Enzyme Immunoassay kit for the quantification of Human SPINK5 in serum, plasma (EDTA, heparin, citrate) and cell culture supernatants.	
Tested Reactivity	Hu	
Tested Application	ELISA	
Target Name	SPINK5	
Conjugation	HRP	
Conjugation Note	Substrate: TMB and read at 450 nm.	
Sensitivity	156 pg/ml	
Sample Type	Serum, plasma (EDTA, heparin, citrate) and cell culture supernatants.	
Standard Range	312 - 20000 pg/ml	
Sample Volume	100 μΙ	
Precision	Intra-Assay CV: 5.0% Inter-Assay CV: 5.8%	

VAKTI: NS: NETS

~ 5 hours

#### **Application Instructions**

Assay Time

**Properties** 

Form 96 well

Storage instruction Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test

reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual

for detail temperatures of the components.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol SPINK5

Gene Full Name serine peptidase inhibitor, Kazal type 5

Background This gene encodes a multidomain serine protease inhibitor that contains 15 potential inhibitory

domains. The inhibitor may play a role in skin and hair morphogenesis and anti-inflammatory and/or antimicrobial protection of mucous epithelia. Mutations may result in Netherton syndrome, a disorder characterized by ichthyosis, defective cornification, and atopy. Multiple transcript variants encoding

different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

Function Serine protease inhibitor, probably important for the anti-inflammatory and/or antimicrobial protection

of mucous epithelia. Contribute to the integrity and protective barrier function of the skin by regulating the activity of defense-activating and desquamation-involved proteases. Inhibits KLK5, it's major target,

in a pH-dependent manner. Inhibits KLK7, KLK14 CASP14, and trypsin. [UniProt]


Highlight Related products:

New ELISA data calculation tool: Simplify the ELISA analysis by GainData

PTM Proteolytically processed by furin in individual domains (D1, D5, D6, D8 through D11, and D9 through

D15) exhibiting various inhibitory potentials for multiple proteases. [UniProt]

Cellular Localization Secreted. [UniProt]



#### ARG82533 Human SPINK5 ELISA Kit standard curve image

ARG82533 Human SPINK5 ELISA Kit results of a typical standard run with optical density reading at 450 nm.