

## ARG82600 Human CD140a / PDGFRA ELISA Kit

Package: 96 wells

Store at: 4°C

### Summary

Product Description	ARG82600 Human CD140a / PDGFRA ELISA Kit is an Enzyme Immunoassay kit for the quantification of Human CD140a / PDGFRA in serum, plasma and cell culture supernatants.
Tested Reactivity	Hu
Tested Application	ELISA
Specificity	<p>This kit could assay both natural and recombinant Human CD140a / PDGFRA.</p> <p>No significant cross-reactivity or interference was observed in the following samples: Human: IFN gamma, IL1 beta, IL2, IL4, IL5, IL6, IL8, IL10, IL12, IL17A, IL18, IL21, IL22, IL23, MCP1, TGF beta 1, TNF alpha and VEGF. Mouse: GM-CSF, IFN gamma, IL1 beta, IL2, IL4, IL6, IL10, IL17A and TNF alpha. Rat: IFN gamma, IL1 beta, IL4, IL6, IL10 and TNF alpha.</p>
Target Name	CD140a / PDGFRA
Conjugation	HRP
Conjugation Note	Substrate: TMB and read at 450 nm.
Sensitivity	31.3 pg/ml
Sample Type	Serum, plasma and cell culture supernatants.
Standard Range	62.5 - 4000 pg/ml
Sample Volume	20 µl
Precision	Intra-Assay CV: 3.1% Inter-Assay CV: 5.2%
Alternate Names	Platelet-derived growth factor receptor alpha; PDGFR2; Alpha-type platelet-derived growth factor receptor; RHEPDGFRA; PDGFR-2; Platelet-derived growth factor receptor 2; CD140A; Platelet-derived growth factor alpha receptor; CD140 antigen-like family member A; CD antigen CD140a; PDGF-R-alpha; Alpha platelet-derived growth factor receptor; CD140a antigen; EC 2.7.10.1; PDGFR-alpha

### Application Instructions


Assay Time	~ 2.5 hours
------------	-------------

### Properties

Form	96 well
Storage instruction	Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual for detail temperatures of the components.
Note	For laboratory research only, not for drug, diagnostic or other use.

### Bioinformation

<b>Gene Symbol</b>	PDGFRA
<b>Gene Full Name</b>	platelet-derived growth factor receptor, alpha polypeptide
<b>Background</b>	This gene encodes a cell surface tyrosine kinase receptor for members of the platelet-derived growth factor family. These growth factors are mitogens for cells of mesenchymal origin. The identity of the growth factor bound to a receptor monomer determines whether the functional receptor is a homodimer or a heterodimer, composed of both platelet-derived growth factor receptor alpha and beta polypeptides. Studies suggest that this gene plays a role in organ development, wound healing, and tumor progression. Mutations in this gene have been associated with idiopathic hypereosinophilic syndrome, somatic and familial gastrointestinal stromal tumors, and a variety of other cancers. [provided by RefSeq, Mar 2012]
<b>Function</b>	Tyrosine-protein kinase that acts as a cell-surface receptor for PDGFA, PDGFB and PDGFC and plays an essential role in the regulation of embryonic development, cell proliferation, survival and chemotaxis. Depending on the context, promotes or inhibits cell proliferation and cell migration. Plays an important role in the differentiation of bone marrow-derived mesenchymal stem cells. Required for normal skeleton development and cephalic closure during embryonic development. Required for normal development of the mucosa lining the gastrointestinal tract, and for recruitment of mesenchymal cells and normal development of intestinal villi. Plays a role in cell migration and chemotaxis in wound healing. Plays a role in platelet activation, secretion of agonists from platelet granules, and in thrombin-induced platelet aggregation. Binding of its cognate ligands - homodimeric PDGFA, homodimeric PDGFB, heterodimers formed by PDGFA and PDGFB or homodimeric PDGFC -leads to the activation of several signaling cascades; the response depends on the nature of the bound ligand and is modulated by the formation of heterodimers between PDGFRA and PDGFRB. Phosphorylates PIK3R1, PLCG1, and PTPN11. Activation of PLCG1 leads to the production of the cellular signaling molecules diacylglycerol and inositol 1,4,5-trisphosphate, mobilization of cytosolic Ca(2+) and the activation of protein kinase C. Phosphorylates PIK3R1, the regulatory subunit of phosphatidylinositol 3-kinase, and thereby mediates activation of the AKT1 signaling pathway. Mediates activation of HRAS and of the MAP kinases MAPK1/ERK2 and/or MAPK3/ERK1. Promotes activation of STAT family members STAT1, STAT3 and STAT5A and/or STAT5B. Receptor signaling is down-regulated by protein phosphatases that dephosphorylate the receptor and its down-stream effectors, and by rapid internalization of the activated receptor. [UniProt]
<b>Highlight</b>	Related products: <a href="#">CD140a antibodies</a> ; <a href="#">CD140a ELISA Kits</a> ; New ELISA data calculation tool: <a href="#">Simplify the ELISA analysis by GainData</a>
<b>PTM</b>	N-glycosylated.  Ubiquitinated, leading to its degradation.  Autophosphorylated on tyrosine residues upon ligand binding. Autophosphorylation occurs in trans, i.e. one subunit of the dimeric receptor phosphorylates tyrosine residues on the other subunit. Phosphorylation at Tyr-731 and Tyr-742 is important for interaction with PIK3R1. Phosphorylation at Tyr-720 and Tyr-754 is important for interaction with PTPN11. Phosphorylation at Tyr-762 is important for interaction with CRK. Phosphorylation at Tyr-572 and Tyr-574 is important for interaction with SRC and SRC family members. Phosphorylation at Tyr-988 and Tyr-1018 is important for interaction with PLCG1. [UniProt]
<b>Cellular Localization</b>	Cell membrane; Single-pass type I membrane protein. Note=The activated receptor is rapidly internalized and degraded. [UniProt]



ARG82600 Human CD140a / PDGFRA ELISA Kit standard curve image

ARG82600 Human CD140a / PDGFRA ELISA Kit results of a typical standard run with optical density reading at 450 nm.