

ARG83089 Human C Reactive Protein ELISA Kit

Package: 96 wells
Store at: 4°C

Summary

Product Description	ARG83089 Human C Reactive Protein ELISA Kit is an Enzyme Immunoassay kit for the quantitative of Human C Reactive Protein in Serum.
Tested Reactivity	Hu
Tested Application	ELISA
Target Name	C Reactive Protein
Conjugation	HRP
Conjugation Note	Substrate: TMB and read at 450 nm.
Sensitivity	10 ng/ml
Sample Type	Serum
Standard Range	100 – 10,000 ng/ml
Sample Volume	20 µl
Precision	Intra-Assay CV: less than 10 % Inter-Assay CV: less than 10 %
Alternate Names	1-205; PTX1; C-reactive protein

Application Instructions

Assay Time	60 min
------------	--------

Properties

Form	96 well
Storage instruction	Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual for detail temperatures of the components.
Note	For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol	CRP
Gene Full Name	C-reactive protein, pentraxin-related
Background	<p>C-Reactive Protein (CRP) is an acute-phase protein, produced exclusively in the liver. Interleukin-6 is the mediator for the synthesis by the hepatocytes of CRP, a pentamer of approximately 120.000 Daltons. CRP is present in the serum of normal persons at concentrations ranging up to 5mg/l. The protein is produced by the fetus and the neonate and it does not pas the placental barrier, as such it can be used for the early detection of neonatal sepsis.</p> <p>Because febrile phenoena, leukocyte count and erythrocyte sedimentation rate (ESR) are often</p>

misleading, investigators and clinicians now prefer a quantitative CRP determination as a marker for acute inflammation and tissue necrosis. Within 6 hours of an acute inflammatory challenge the CRP level starts to rise. Serum concentration of CRP increases significantly in cases of both infectious and non-infectious inflammation, of tissue damage and necrosis and in the presence of malignant tumours. CRP is present in the active stages of inflammatory disorders like rheumatoid arthritis, ankylosing spondylitis, Reiter's syndrome, psoriatic arthropathy, systemic lupus erythematosus, polyarteritis, ulcerative colitis and Crohn's disease. Injuries causing tissue breakdown and necrosis are associated with increases in serum CRP which are seen in thermal burns, major surgery and myocardial infarction. Widespread malignant disease with carcinoma of the lung, stomach, colon, breast, prostate and pancreas, Hodgkin's disease, non-Hodgkin's lymphoma and lymphosarcoma will give rise to high levels of CRP resulting from tissue damage by invading tumour cells. CRP, therefore may be used to monitor malignancy. The CRP-level increases dramatically following microbial infections, and this may be particularly helpful for the diagnosis and monitoring of bacterial septicemia in neonates and other immunocompromised patients at risk. In children, CRP is useful for differential diagnosis of bacterial and viral meningitis. Because the biological half-life of this protein is only 24 hours, CRP accurately parallels the activity of the inflammation process and the CRP concentration decreases much faster than ESR^{1,2} or any other acute phase parameter, which is particularly useful in monitoring appropriate treatment of bacterial diseases with antibiotics. C-Reactive Protein measurements during the early and late post transplant period of bone marrow and organ transplantations is particularly useful in the management of interfering infections in these immunosuppressed patients.

Highlight

Related products:

[CRP antibodies](#); [CRP ELISA Kits](#); [CRP Duos / Panels](#);


New ELISA data calculation tool:

[Simplify the ELISA analysis by GainData](#)

Research Area

Cell Biology and Cellular Response kit; Cell Death kit; Gene Regulation kit; Immune System kit; Metabolism kit

Images



ARG83089 Human C Reactive Protein ELISA Kit standard curve image

ARG83089 Human C Reactive Protein ELISA Kit results of standard run with optical density reading at 450 nm