

Product datasheet

info@arigobio.com

ARG83344 Human PNP / Purine nucleoside phosphorylase ELISA Kit

Package: 96 wells Store at: 4°C

Summary

Product Description ARG83344 Human PNP / Purine nucleoside phosphorylase ELISA Kit is an Enzyme Immunoassay kit for

the quantification of Human PNP / Purine nucleoside phosphorylase in serum, plasma and cell culture

supernatants.

Tested Reactivity Hu

Tested Application ELISA

Specificity There is no detectable cross-reactivity with other relevant proteins.

Target Name PNP / Purine nucleoside phosphorylase

Conjugation HRP

Conjugation Note Substrate: TMB and read at 450 nm.

Sensitivity 15 pg/ml

Detection Range 31.2 pg/ml - 2000 pg/ml

Sample Type Serum, Plasma and Cell culture supernatants.

Alternate Names EC 2.4.2.1; Purine nucleoside phosphorylase; Inosine-guanosine phosphorylase; PNP; PRO1837; Inosine

phosphorylase; PUNP; NP

Application Instructions

Assay Time ~ 5 hours

Properties

Form 96 well

Storage instruction Store the kit at 2-8°C. Keep microplate wells sealed in a dry bag with desiccants. Do not expose test

reagents to heat, sun or strong light during storage and usage. Please refer to the product user manual

for detail temperatures of the components.

Note For laboratory research only, not for drug, diagnostic or other use.

Bioinformation

Gene Symbol PNP

Gene Full Name purine nucleoside phosphorylase

Background This gene encodes an enzyme which reversibly catalyzes the phosphorolysis of purine nucleosides. The

enzyme is trimeric, containing three identical subunits. Mutations which result in nucleoside

phosphorylase deficiency result in defective T-cell (cell-mediated) immunity but can also affect B-cell immunity and antibody responses. Neurologic disorders may also be apparent in patients with immune defects. A known polymorphism at aa position 51 that does not affect enzyme activity has been described. A pseudogene has been identified on chromosome 2. [provided by RefSeq, Jul 2008]

www.arigobio.com arigo.nuts about antibodies 1/2

Function The purine nucleoside phosphorylases catalyze the phosphorolytic breakdown of the N-glycosidic bond


in the beta-(deoxy)ribonucleoside molecules, with the formation of the corresponding free purine

bases and pentose-1-phosphate. [UniProt]

Research Area Neuroscience antibody

Cellular Localization Cytoplasm, cytoskeleton. Cytoplasm

Images



ARG83344 Human PNP / Purine nucleoside phosphorylase ELISA Kit standard curve image

ARG83344 Human PNP / Purine nucleoside phosphorylase ELISA Kit results of a typical standard run with optical density reading at 450 nm.